Europäisches Patentamt

European Patent Office

Office européen des brevets

(11) EP 0 982 300 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication: 01.03.2000 Bulletin 2000/09

(21) Application number: 98403351.4

(22) Date of filing: 31.12.1998

(51) Int. Cl.⁷: **C07D 295/088**, C07C 211/08, C07D 211/04, C07D 295/185, C07D 211/62, C07D 211/70, C07D 207/20, A61P 25/28

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 29.07.1998 EP 98401944

(71) Applicant: SOCIETE CIVILE BIOPROJET F-75003 Paris (FR)

(72) Inventors:

- Schwartz, Jean-Charles
 75014 Paris (FR)
- Arrang, Jean-Michel 91410 Dourdan (FR)
- Garbarg, Monique 75017 Paris (FR)

- Lecomte, Jeanne-Marie 75003 Paris (FR)
- Ligneau, Xavier 75013 Paris (FR)
- Schunack, Walter G. 14129 Berlin (DE)
- Stark, Holger
 14199 Berlin (DE)
- Ganellin, Charon Robin Welwyn, Hert AL6 0TD (GB)
- Leurquin, Fabien
 London E2 6DZ (GB)
- Sigurd, Elz
 12107 Berlin (DE)

(74) Representative:

Le Guen, Gérard et al CABINET LAVOIX 2, place d'Estienne d'Orves 75441 Paris Cédex 09 (FR)

- (54) Non-imidazole alkylamines as histamine H3 receptor ligands and their therapeutic applications
- (57) Compounds of formula (A):

W]-N $\stackrel{R^1}{\underset{R^2}{\sim}}$ (A)

a N-substituted piperazino group as defined.

Compounds (A) are useful for preparing medicaments acting as antagonists and/or agonists at the H₃-receptors of histamine.

wherein:

- W is a residue which imparts antagonistic and/or agonistic activity at histamine H₃-receptors when attached to an imidazole ring in 4(5) position.
- R¹ and R² may be identical or different and represent each independently
 - a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
 - a saturated nitrogen-containing ring (i) as defined,
 - a non-aromatic unsaturated nitrogen-containing ring (ii) as defined,
 - a morpholino group, or

Description

10

[0001] The present invention relates to alkylamines of formula (A) as defined hereafter, to their preparation and to their therapeutic applications.

[0002] Antagonists of histamine H₃-receptor are known especially to increase synthesis and release of cerebral histamine. Through this mechanism, they induce an extended wakefullness, an improvement in cognitive processes, a reduction in food intake and a normalization of vestibular reflexes (Schwartz et al., Physiol. Rev., 1991, 71: 1-51).

[0003] Whence these agents are potentially useful in several central nervous system disorders such as Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness

[0004] Histamine H₃-receptor agonists are known to inhibit the release of several neurotransmitters including histamine, monoamines and neuropeptides and thereby exert sedative and sleep-promoting effects in brain. In peripheral tissues, H₃-receptor agonists exert namely anti-inflammatory, anti-nociceptive, gastro-intestinal, antisecretory smooth muscle decontracting activities.

[0005] All the H₃ receptor antagonist or agonist compounds known so far resemble histamine in possessing an imidazole ring generally monosubstituted in 4(5)-position (Ganellin et al., Ars Pharmaceutica, 1995, 36:3, 455-468; Stark et al., Drug of the Future, 1996, 21(5), 507-520).

[0006] Numerous patents and patent applications are directed to antagonist and/or agonist compounds having such structure, in particular EP 197 840, EP 494 010, WO 93/14070, WO 96/29315, WO 92/15 567, WO 93/20061, WO 93/20062, WO 95/11894, US 5 486 526, WO 93/12107, WO 93/12108, WO 95/14007, WO 95/06037, WO 97/29092, EP 680 960, WO 96/38141, WO 96/38142, WO 96/40126.

[0007] In the litterature, Plazzi et al., Eur. J. Med. Chem. 1995, 30, 881, Clitherow et al., Bioorg. & Med. Chem. Lett. 6 (7), 833-838 (1996) Wolin et al., Bioorg. & Med. Chem. Lett; 8, 2157 (1998) can be cited also in this respect.

[0008] Nevertheless, such imidazole derivatives may show drawbacks such as poor blood-brain barrier penetration, interaction with cytochrome P-450 proteins and/or some hepatic and ocular toxicities.

[0009] Non-imidazole known neuro-active compounds such as betahistine (J-M. Arrang et al., Eur. J. Pharmacol. 1985, 111: 72-84), phencyclidine (J-M. Arrang et al., Eur. J. Pharmacol. 1988, 157: 31-35), dimaprit (J-C Schwartz et al., Agents Actions 1990, 30: 13-23), clozapine (M. Kathmann et al., Psychopharmacology 1994, 116: 464-468), and sesquiterpenes (M. Takigawa et al., JP 06 345 642 (20 Dec 1994)) were suggested to display H₃-receptor antagonism but all these compounds have only very low potency.

[0010] These compounds were previously known as therapeutic agent before the discovery and characterization of the histamine H₃-receptor, in particular as neuro-active agents for example as neuroleptic (clozapine) or psychotomimetic (Phencyclidine) agent.

[0011] When tested at the H₃-receptor, these compounds were shown to display much lower potency than the imidazole-containing compounds described in patent applications quoted above.

[0012] Attempts at replacing the imidazole ring was generally not successful and no potent H₃-receptor ligands not containing such ring was reported in the literature up to now.

[0013] These investigations showed the importance of the 4(5)-imidazole moiety.

[0014] The objective of the invention is to provide new potent H_3 -receptor ligands which may reduce the above-mentioned drawbacks.

[0015] The present invention provides new compounds, the structure of which does not contain an imidazole moiety, which are useful as histamine H₃-receptor ligands.

[0016] The compounds of the invention have the following general formula (A):

 $[W] - N <_{R^2}^{R^1}$ (A)

50 in which:

45

55

- W is a residue which imparts antagonistic and/or agonistic activity at histamine H₃-receptors when attached to an imidazole ring in 4(5)-position;
- R¹ and R² may be identical or different and represent each independently

• a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,

· a saturated nitrogen-containing ring

with m ranging from 2 to 8, or

5

10


15

20

25

30

· a non-aromatic unsaturated nitrogen-containing ring



with p and q being from 0 to 3 independently and r being from 0 to 4, provided that p and q are not simulteously 0 and $2 \le p + q + r \le 8$, R^{a-d} being independently a hydrogen atom or a lower alkyl, cycloalkyl, or carboalkoxy group, or

- · a morpholino group, or
- a N-substituted piperazino group:

with R being a lower alkyl, cycloalkyl, carboalkoxy, aryl, arylalkyl, an alkanoyl or aroyl group.

[0017] The inventors have found, surprisingly, that antagonist and/or agonist compounds can be obtained by substituting a di(alkyl) or (cycloalkyl)amine, or a non-aromatic nitrogen-containing ring —NR¹R² as above-defined for the imidazole ring, in known antagonist and/or agonist imidazole derivatives.

[0018] It is also believed that antagonist and/or agonist activity can be foreseen, by equivalence, for compounds according to formula (A) having a W residue of imidazole derivatives which were suggested in the prior art as H₃ antagonists or agonists, and further for those W residues which would belong to future imidazole derivatives having substantial H₃ antagonist and/or agonist activity.

[0019] Moreover, the inventors have observed that such non-imidazole analogues can provide potent antagonist and/or agonist activity.

[0020] In this regards, they have prepared novel non-imidazole alkylamines analogues of formula (A) corresponding to known imidazole derivatives in particular from the above-mentioned prior art.

[0021] The invention also relates to the addition salts which the compounds form with pharmaceutically acceptable acids. The pharmaceutically acceptable salts comprise the nontoxic salt of inorganic or organic acids. Examples of these salts include the hydrochloride, the hydrobromide or the hydrogen maleate or hydrogen oxalate.

[0022] The present invention also encompasses the hydrates of the compounds, the hydrated salts of these compounds and the polymorphic crystalline structures.

[0023] When the compounds can exist in one or a number of isomeric forms according to the number of asymmetric centres in the molecule, the invention relates both to all the optical isomers and to their racemic modifications and the corresponding diastereoisomers. The separation of the diastereoisomers and/or of the optical isomers can be carried out according to methods known per se.

[0024] The present invention also encompasses all the possible tautomeric forms of the compounds, whether these tautomers occur in isolated form or in the form of mixtures.

[0025] According to the invention, lower alkyl or cycloalkyl is intended to mean a linear or branched alkyl group containing from 1 to 6 carbon atoms, or a saturated carbocycle containing 3 to 6 carbon atoms.

[0026] Typically examples of lower alkyl are methyl, ethyl, propyl, isopropyl and butyl groups.

[0027] A preferred group of compounds according to the invention comprises those with R1 and R2 representing inde-

pendently a lower alkyl group, especially an ethyl group.

[0028] Preferred compounds are also those of formula (A) in which R¹ and R² taken together with the nitrogen atom to which they are attached, form a saturated nitrogen-containing ring:

i) N (CR^aR^b)_m

especially with m being 4, 5 or 6, optionally substituted with an alkyl group (Ra), preferably a methyl group.

[0029] The groups Ra and Rb are identical or different for each (CRaRb) moiety.

[0030] Piperidyl and pyrrolidinyl moieties are especially preferred.

[0031] Another preferred group of compounds comprises compounds (A) in which R¹ and R² taken together with the nitrogen atom to which they are attached, form a non-aromatic unsaturated nitrogen-containing ring:

ii) N (CHR^d)_q+CR^c

25

20

5

10

especially with p, q, and r being independently 1 or 2.

[0032] In this group, more preferred compounds are those with p being 2 and q and r each being 1.

[0033] A sub-class in this group comprises compounds with Ra-d being each a hydrogen atom.

[0034] When NR¹R² is a nitrogen-containing ring i) or ii) as above-defined, the latter is preferably substituted with one or two lower alkyl group(s), especially a methyl group.

[0035] The position for substitution is preferably selected according the following order:

meta>para>ortho.

35 [0036] The position for substitution is preferably selected according the following order:

meta>para>ortho.

[0037] In this group, for nitrogen-containing ring bearing only one substituent, this latter is preferably in meta position with respect to the nitrogen-atom.

[0038] For nitrogen-containing ring bearing two substituents, meta-meta substitution is preferred, especially when these two substituents are in trans-relation.

[0039] According to the invention, piperidyl or pyrrolidinyl moiety substituted in meta or meta-meta position, especially with a methyl group, give particularly preferred compounds.

[5 [0040] When NR¹R² represents a N-substituted piperazino group, R may be a lower alkyl e.g. methyl.

[0041] Typical examples of group R being an aryl or arylalkyl moiety are phenyl and benzyl.

[0042] R may be also an alkanoyl or aroyl group e.g. acetyl or benzoyl.

[0043] In all the possible groups for R, the alkyl moiety refers to a linear or branched chain containing from 1 to 6 carbon atoms.

[0044] The cycloalkyl group refers to a saturated carbocycle containing 3 to 7 carbon atoms.

[0045] When R represents an aryl or arylalkyl group, the aryl moiety is especially a phenyl group optionally substituted with one or more substituents selected from halogen atoms, advantageously selected from fluorine, chlorine and bromine, or a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino, α-hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group.

[0046] R may be also an optionally substituted benzoyl, the substituent being as defined above with reference to the phenyl group.

[0047] Typical example of —NR¹R² representing a N-substituted piperazino group is N-acetylpiperazino.

[0048] According to one aspect, the compounds of the invention have the following general formula (I):

$$(R^3)_{n3}$$
 $+ C_n H_{2n} - N$ $+ N_{R^2}$ (I)

in which:

5

10

20

- C_nH₂n is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- 5 n₃ is an integer from 0 to 5;
 - R³ represents each independently
 - · a halogen atom,
 - a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino, α-hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
 - or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated
 or unsaturated ring or a benzene ring.
- \sim R¹ and R² are as above-defined in formula (A).

[0049] A preferred group of compounds according to the invention is the group composed of compounds of formula (I) in which X is an oxygen atom.

[0050] Another preferred group of compounds comprises compounds (I) in which -C_nH_{2n}- is a linear chain -(CH₂)_n-with n being as previously defined.

[0051] Preferred compounds are also those with n varying from 3 to 5, and with n being more preferably 3.

[0052] A sub-class of compounds according to the invention comprises the compounds of formula (I) with n₃ being zero that is those having an unsubstituted phenyl molety.

[0053] Another group of compounds according to the invention is composed of compounds containing one or more substituents R³ which may be identical or different. In this group, the compounds having a mono- or di-substituted (n₃ = 1 or 2) phenyl moiety are preferred and those mono-substituted with one group R³ as defined above in para-position are particularly preferred.

[0054] Among these compounds, (n_3 being 1) R^3 is preferably a halogen atom or a cyano, nitro, alkanoyl, alkyloximino or α -hydroxyalkyl group.

[0 [0055] Still more preferred compounds are those with R³ being CN, NO₂, COCH₃, COC₂H₅, H₃C-C=N-OH, H₃C-CH-OH.

[0056] R³ being a halogen atom may be advantageously selected from fluorine, chlorine and bromine.

[0057] R³ being an aryl group, may be especially a phenyl group.

[0058] In the other substituents R³, the aryl moiety is advantageously a phenyl moiety.

[0059] R³ being an aryloxy group may be especially a phenoxy group.

[0060] According to the invention, alkanoyl is intended to mean a group containing an alkyl moiety as defined above.

[0061] Typical examples of R³ being an alkanoyl, aroyl or arylalkanoyl group are acetyl, butyryl and propionyl groups, benzoyl group or phenylacetyl group.

[0062] Typical examples of R³ forming together with the carbon atoms of the phenyl ring to which it is fused, a saturated ring leads to 5,6,7,8-tetrahydronaphthyl or forming a benzene ring leads to a naphthyl moiety.

[0063] According to the invention, alkenyl or alkynyl group may contain advantageously from 1 to 8 carbon atoms, in particular from 1 to 6 carbon atoms and preferably 1 to 4 carbon atoms.

[0064] In carboalkoxy, carboxyamido or carboxamide groups, the hydrocarbon chain is saturated, linear or branched and contains an alkyl moiety as defined above.

5 [0065] In alkoxy, alkyloximino, arylalkyl or α-hydroxyalkyl group, the alkyl moiety is as previously defined also.

[0066] Particularly preferred compounds are:

1-(5-phenoxypentyl)-piperidine

	1-(5-phenoxypentyl)-pyrrolidine
	N-methyl-N-(5-phenoxypentyl)-ethylamine
	1-(5-phenoxypentyl)-morpholine
	N-(5-phenoxypentyl)-hexamethyleneimine
5	N-ethyl-N-(5-phenoxypentyl)-propylamine
	1-(5-phenoxypentyl)-2-methyl-piperidine
	1-(5-phenoxypentyl)-4-propyl-piperidine
	1-(5-phenoxypentyl)-4-methyl-piperidine
	1-(5-phenoxypentyl)-3-methyl-piperidine
10	1-acetyl-4-(5-phenoxypentyl)-piperazine
	1-(5-phenoxypentyl)-3,5-trans-dimethyl-piperidine
	1-(5-phenoxypentyl)-3,5-cis-dimethyl-piperidine
	1-(5-phenoxypentyl)-2,6-cis-dimethyl-piperidine
	4-carboethoxy-1-(5-phenoxypentyl)-piperidine
15	3-carboethoxy-1-(5-phenoxypentyl)-piperidine
	1-(5-phenoxypentyl)-1,2,3,6-tetrahydropyridine
	1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
20	1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
	1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine
	1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine
	1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
25	1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
	1-{5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
	1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
	1-(5-phenoxypentyl)-2,5-dihydropyrrole
	1-{5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
30	1-(4-phenoxybutyl)-pyrrolidine
	1-(6-phenoxyhexyl)-pyrrolidine
	1-(5-phenylthiopentyl)-pyrrolidine
	1-(4-phenylthiobutyl)-pyrrolidine
	1-(3-phenoxypropyl)-pyrrolidine
35	1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
	1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
40	1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
,,,	N-[3-(4-nitrophenoxy)-propyl]-diethylamine
	N-[3-(4-cyanophenoxy)-propyl]-diethylamine
	1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
	1-{5-[4-(phenylacetyl)-phenoxy]-pentyl}-pyrrolidine
45	N-[3-(4-acetylphenoxy)-propyl]-diethylamine
	1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
	1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
50	1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
-	1-[5-(4-cyanophenoxy)-pentyl]-piperidine
	N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
	N-[2-(4-cyanophenoxy)-ethyl]-diethylamine
	N-[3-(4-cyanophenoxy)-propyl]-dimethylamine
55	N-[4-(4-cyanophenoxy)-butyl]-diethylamine
<i>33</i>	N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
	1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
	1-[3-(4-cyanophenoxy)-propyl]-piperidine
	i-[o-(+-cyanophenoxy/-phopyi]-phendine

- N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
- N-[6-(4-cyanophenoxy)-hexyl]-diethylamine
- N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
- N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
- 4-(3-diethylaminopropoxy)-acetophenone-oxime
 - 1-[3-(4-acetylphenoxy)-propyl]-piperidine
 - 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 - 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine
 - 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
- 10 1-[3-(4-propionylphenoxy)-propyl]-piperidine
 - 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
 - 1-[3-(4-formylphenoxy)-propyl]-piperidine
 - 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
 - N-[3-(4-propionylphenoxy)-propyl]-diethylamine
 - 1-[3-(4-butyrylphenoxy)-propyl]-piperidine
 - 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine

[0067] More preferred compounds are:

- 20 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 - N-[3-(4-cyanophenoxy)-propyl]-diethylamine
 - N-[3-(4-acetylphenoxy)-propyl]-diethylamine
 - 1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
 - N-[4-(4-cyanophenoxy)-butyl]-diethylamine
- 25 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 - N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
 - N-3-[4-(1-hydroxyethyl)-phenoxyl-propyl-diethylamine
 - 4-(3-diethylaminopropoxy)-acetophenone-oxime
 - 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
- 30 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 - 1-[3-(4-propionylphenoxy)-propyl]-piperidine

[0068] Compounds of formula (I) in which:

- -NR¹R² is a pyrrolidinyl group, C_nH_{2n} is a linear chain -(CH₂)_n- and n₃ is zero, X being an oxygen atom with n ranging from 3 to 5, or X being a sulfur atom with n being 4 or 5;
 - -NR¹R² is a piperidinyl group, C_nH_{2n} is a linear chain -(CH₂)_n- and X is an oxygen atom, n₃ being zero with n being 2, 5 or 8 or n₃ being 1 with R³ being 4-CN and n being 5;
 - -NR¹R² is a diethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain -(CH₂)_n- and n₃ is 1, R³ being 4-NO₂ or 4-COCH₃ with n being 3 or R³ being 4-CN with n being 2 to 4;
 - -NR¹R² is a dimethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain -(CH₂)_n- and n³ is 1, R³ being 4-CN with n being 3,

are known in the art.

- [0069] A subject of the invention is thus the use of these compounds as antagonists at the histamine H₃-receptors, in particular to prepare medicaments acting as H₃-antagonists intended for the treatments detailed below.
 - [0070] According to a second aspect, the object of the present invention is non-imidazole compounds analogous to the compounds disclosed in WO 96/29315 and WO 93/14070.
- [0071] Thus, a first sub-class of the compounds (A) of the invention is defined by the compounds having the following general formula (IIa) and (IIb):

40

5

$$\begin{array}{c|c} R^{1} & \text{N-(chain A II)} - X^{II} - (\text{chain B II)} - Y^{II} & \text{(IIa)} \end{array}$$

or

$$\begin{array}{c}
R^1 \longrightarrow N \longrightarrow (\text{chain A}^{||}) \longrightarrow X^{||} \longrightarrow Y^{||} \\
R^2 \longrightarrow N \longrightarrow (\text{chain A}^{||}) \longrightarrow X^{||} \longrightarrow Y^{||} \\
\end{array} (||b|)$$

in which

5

10

20

25

55

- 15 R¹ and R² are as defined with reference to general formula (A);
 - the chain A^{II} represents a saturated or unsaturated, straight or branched hydrocarbon chain containing 1 to 6 carbon atoms, it being possible for the saturated hydrocarbon chain to be interrupted by a hetero atom such as a sulphur atom;
 - X^{II} represents an oxygen or sulphur atom, -NH-, -NHCO-, -N(alkyl)CO-, -NHCONH-, -NH-CS-NH-, -NHCS-, -O-CO-, -CO-O-, -OCONH-, -OCON(alkyl)-, -OCONH-CO-, -CONH-, -CON(alkyl)-, -SO-, -CO-, -CHOH- or -NR_{II}-C(=NR"_{II})-NR'_{II}-, R_{II} and R'_{II} denoting a hydrogen atom or a lower alkyl radical and R"_{II} a hydrogen atom or another powerful electronegative group, such as a cyano or COY₁^{II} group, Y₁^{II} denoting an alkoxy group;
 - the chain B^{II} represents a straight alkylene chain -(CH₂)_{nII}-, n being an integer which can vary between 1 and 5 or a branched alkylene chain containing from 2 to 8 carbon atoms, the alkylene chain being optionally interrupted by one or a number of oxygen or sulphur atoms, or a group -(CH₂)_{nII}-O- or -(CH₂)_{nII}-S- where n_{II} is an integer equal to 1 or 2;

[0072] Y^{II} represents a straight or branched alkyl group containing 1 to 8 carbon atoms; a cycloalkyl containing 3 to 6 carbon atoms; a bicycloalkyl group; a cycloalkenyl group; an aryl group such as an optionally substituted phenyl group; a 5- or 6-membered heterocyclic radical containing one or two heteroatoms chosen from nitrogen and sulphur atoms, the said heterocyclic radical optionally being substituted; or also a bicyclic radical resulting from the fusion of a benzene ring to a heterocycle as defined above.

[0073] The chain A can be a straight alkylene chain - $(CH_2)_{n|l}$ -, $n_{|l}$ representing an integer between 1 and 6 carbon atoms, preferably between 1 and 4 carbon atoms, or a branched alkylene chain, preferably a chain substituted by one or a number of methyl or ethyl radicals.

[0074] The chain A^{II} can also be a straight or branched unsaturated alkylene chain, and can be, for example, the allyl group.

[0075] When Y^{II} represents a cyclcoalkyl group, the latter can be, for example, cyclopentyl, cyclohexyl or a bicycloalkyl group.

[0076] When Y^{II} represents a substituted phenyl group, the phenyl group can be mono- or polysubstituted, for example, by a halogen, by a lower alkyl, for example CH₃, by CF₃, CN, COCH₃, COOR^{II}₁ or OR^{II}₁, R^{II}₁ representing a lower alkyl, for example COOCH₃, the NO₂ group or the group NR^{II}₂R^{II}₃, R^{II}₂ and R^{II}₃ representing a hydrogen atom and/or a lower alkyl radical ("lower alkyl" means an alkyl radical containing at most 6 carbon atoms).

[0077] When Y^{II} represents a heterocyclic radical, the latter can be, for example, the pyridyl radical, the pyridyl N-oxide radical or the pyrazinyl radical, optionally mono- or polysubstituted by NO₂, CF₃, CH₃, NH₂, a halogen such as CI, the COOCH₃ group or also the thiazolyl radical.

[0078] When Y^{II} represents a bicyclic radical resulting from the fusion of a benzene ring to a heterocycle, the radical can be, for example, the benzothiazolyl radical.

[0079] A second sub-class of the compounds (A) according to the invention comprises the compounds having the above-formulae (IIa) and (IIb) in which:

- R¹R² are as defined with reference to general formula (A);
- the chain A" represents an unbranched, branched or unsaturated alkyl group -(CH₂)_{n|l}- where n_{|l} is an integer which can vary between 1 and 8 and preferably between 1 and 4; an unbranched or branched alkene group comprising from 1 to 8 carbon atoms and preferably 1 to 4 carbon atoms; an unbranched or branched alkyne group comprising from 1 to 4 carbon atoms;
- the group X^{II} represents -OCONH-; -OCON(alkyI)-; -OCON(alkene)-; -OCO-; -OCSNH-; -CH₂-; -O-; -OCH₂CO-; -S-; -CO-; -CS-; amine; alkene;

— the chain B^{II} represents an unbranched, branched or unsaturated lower alkyl comprising from 1 to 8 carbon atoms and preferably 1 to 5 carbon atoms; -(CH₂)_{nII}(hetero atom)- where the hetero atom is preferably a sulphur or oxygen atom; n_{II} being an integer which can vary between 1 and 5, preferably between 1 and 4;

5

10

15

20

the group YII represents a phenyl group, unsubstituted or mono- or polysubstituted with one or more identical or different substituents selected from halogen atoms, OCF3, CHO, CF3, SO2N(alkyl)2 such as SO2N(CH3)2, NO2, S(alkyl), S(aryl), SCH2(phenyl), an unbranched or branched alkene, an unbranched or branched alkyne optionally substituted with a trialkylsilyl radical, -O(alkyl), -O(aryl), -CH2CN, a ketone, an aldehyde, a sulphone, an acetal, an alcohol, a lower alkyl, -CH=CH-CHO, -C(alkyl)=N-OH, -C(alkyl)=N-O(alkyl) and other keto derivatives, -CH=NOH, -CH=NO(alkyl), and other aldehyde derivatives, -C(alkyl)=NH-NH-CONH2, an O-phenyl or -OCH2(phenyl) group, -C(cycloalkyl)=NOH, -C(cycloalkyl)=N-O(alkyl), an optionally substituted heterocycle; a heterocycle comprising a sulphur hetero atom; a cycloalkyl; a bicyclic group and preferably a norbornyl group; a phenyl ring fused to a heterocycle comprising a nitrogen hetero atom or to a carbocycle or a heterocycle bearing a keto function; an unbranched or branched lower alkyl comprising from 1 to 8 carbon atoms; an unbranched or branched alkyne comprising from 1 to 8 carbon atoms and preferably 1 to 5 carbon atoms; a linear or branched alkyl mono- or polysubstituted with phenyl groups which are either unsubstituted or mono- or polysubstituted; a phenyl alkyl ketone in which the alkyl group is branched or unbranched or cyclic; a substituted or unsubstituted benzophenone; a substituted or unsubstituted, unbranched or branched or cyclic phenyl alcohol; an unbranched or branched alkene; a piperidyl group; a phenylcycloalkyl group; a polycyclic group, in particular a fluorenyl group, a naphthyl or polyhydronaphthyl group or an indanyl group; a phenol group; a ketone or keto derivative; a diphenyl group; a phenoxyphenyl group; a benzyloxyphenyl group.

[0080] According to the invention, group X^{II} representing an amine is understood to mean a secondary or tertiary amine.

[0081] The alkyl, alkene, alkyne, keto, aldehyde, cycloalkyl, S-alkyl, O-alkyl, phenyl alcohol and phenyl-cycloalkyl groups mentioned above as well as in the remainder of the description and the claims of the present patent comprise from 1 to 8 carbon atoms, and preferably 1 to 5.

[0082] Likewise, keto derivatives are understood to mean any oxime, alkyloxime, hydrazone, acetal, aminal, ketal, thione, carbazone or semicarbazone group and the thio analogues of these derivatives.

[0083] Likewise, by mono- or polysubstituted phenyl and/or benzophenone groups, it is understood to mean that these groups are substituted with one or more identical or different substituents selected from halogen atoms, OCF₃, CHO, CF₃, SO₂N(alkyl)₂, SO₂N(CH₃)₂, NO₂, S(alkyl), S(aryl), SCH₂(phenyl), an unbranched or branched alkene, an unbranched or branched alkyne optionally substituted with a trialkylsilyl radical, -O(alkyl), -O(aryl), -CH₂CN, a ketone, an aldehyde, a sulphone, an acetal, an alcohol, a lower alkyl, -CH=CH-CHO, -C(alkyl)=N-OH, -C(alkyl)=N-O(alkyl) an other keto derivatives, -CH=NOH, -CH=NO(alkyl), and other aldehyde derivatives, -C(alkyl)=NH-NH-CONH₂, an Ophenyl or -OCH₂(phenyl) group, -C(cycloalkyl)=NOH, -C(cycloalkyl)=N-O(alkyl), an optionally substituted heterocycle. [0084] The keto substituent is preferably selected from a linear- or branched-chain aliphatic ketone, it being possible for the said chain to comprise from 1 to 8 carbon atoms and optionally to bear a hydroxyl group, a cycloalkyl ketone, an aryl alkyl ketone or aryl alkenyl ketone in which the aryl group is unsubstituted or mono- or polysubstituted, or a heteroaryl ketone in which the heteroaryl unit is preferably monocyclic.

0 [0085] The acetal substituent preferably consists of an aliphatic acetal comprising from 1 to 8 carbon atoms and optionally bearing a hydroxyl radical.

[0086] Group Y^{II} representing a ketone is understood to mean, in particular, a ketone substituted with an alkyl or aryl group, it being possible for these groups to be substituted or unsubstituted.

[0087] As regards the heterocycles, these comprise from 1 to 3 hetero atoms, preferably sulphur, oxygen or nitrogen atoms.

[0088] The heterocycle substituent is preferably selected from an oxadiazole or an imidazole.

[0089] Preferred compounds (IIa) and (IIb) are those in which X^{II} is selected from -O-, -NH-, -CH₂-, -OCONH-, -NHCO-, -NHCONH-. X^{II} represents more preferably an oxygen atom.

[0090] Preferred compounds (IIa) and (IIb) are also those in which Y^{II} is selected from a linear or branched alkyl group as above defined; a cycloalkyl group as above-defined, in particular cyclopentyl or cyclohexyl group; a phenyl group unsubstituted or mono-substituted, preferred substituent being halogen atom, in particular chorine; a heterocyclic radical, in particular pyridyl N-oxide or pyrazinyl radicals; a bicyclic radical such as a benzothiazolyl radical.

[0091] Y^{II} is preferably a phenyl group at least mono-substituted with -CHO, a ketone, an aldehyde, -CH=CH-CHO, -C(alkyl)=N-OH, -C(alkyl)=N-O(alkyl) and other keto derivatives, -CH=N-OH, -CH=NO(alkyl) and other aldehyde derivatives, -C(cycloalkyl)=NOH, -C(cycloalkyl)=N-O(alkyl).

[0092] According to the invention, Y^{II} represents especially a phenyl group at least mono-substituted with a keto-substituent or an oxime-substituent, or an halogen atom.

[0093] Particularly preferred keto-substituent is cycloalkylketone.

[0094] Other preferred compounds are those wherein Y^{II} represents a phenyl group fused to a carbocycle bearing a keto-function.

[0095] Yet other prefered Y^{II} are phenylalkyl ketone in which the alkyl group is branched or unbranched or cyclic; an optionally substituted benzophenone, a ketone.

[0096] Particularly preferred group YII are a phenyl group unsubstituted or mono-substituted as above-defined.

[0097] The chain A^{II} is preferably a chain -(CH₂)_n^{II}- with n_{II} varying from 1 to 6, preferably from 1 to 4. The chain A^{II} represents especially -(CH₂)₃-.

[0098] Preferred chain B^{II} is -(CH₂)₂- or -(CH₂)₃-.

[0099] Among compounds (IIa) and (IIb), particularly preferred compounds are those in which X^{II} is an oxygen atom, the chain A^{II} represents -(CH₂)₃- and, for compounds of formula (IIa), the chain B^{II} represents -(CH₂)₃- also.

[0100] In this group, YII is preferably an aryl group.

[0101] Preferred group R¹ and R² are as above-defined with reference to formula (A).

[0102] Examples of compounds (IIa) and (IIb) are:

- 15 3,3-Dimethylbutyl 3-piperidinopropyl ether
 - 3-Phenylpropyl 3-piperidinopropyl ether
 - 3-(4-Chlorophenyl)propyl 3-piperidinopropyl ether
 - 2-Benzothiazolyl 3-piperidinopropyl ether
 - N-Phenyl-3-piperidinopropyl carbamate
- 20 N-Pentyl-3-piperidinopropyl carbamate
 - (S)-(+)-N-[2-(3,3-Dimethyl)butyl]-3-piperidinopropyl carbamate
 - 3-Cyclopentyl-N-(3-(1-pyrrolidinyl)propyl)propanamide
 - N-Cyclohexyl-N'-(1-pyrrolidinyl-3-propyl)urea
 - 2-((2-Piperidinoethyl)amino)benzothiazole
- 25 5-Piperidinopentylamine
 - 2-Nitro-5-(6-piperidinohexyl)pyridine
 - 3-Nitro-2-(6-piperidinohexylamino)pyridine
 - 2-(6-Piperidinohexylamino)pyrimidine
 - N-(6-Phenylhexyl)piperidine

[0103] N-phenyl-N'-(N-diethylamino-3-propyl)urea and N-benzyl-N'-(3-piperidinopropyl)guanidine (named compounds 98 and 99 respectively) are also illustrative of compounds (IIa) and (IIb).

[0104] According to a third aspect, the object of the present invention is non-imidazole compounds analogous to the compounds disclosed in EP 197 840.

35 [0105] Thus, a sub-class of compounds (A) according to the invention comprises compounds having the following formula (III)

$$\mathbb{R}^{|||_{3}} \mathbb{N} \mathbb{R}^{1}$$

$$\mathbb{R}^{2}$$

$$(|||)$$

45 in which:

30

40

- NR¹R² is either in 3-position or in 4-position on the piperidyl moiety, R¹ and R² being as defined with reference to formula (A);
- R₂^{III} denotes a linear or branched alkyl group having 1 to 6 carbon atoms; a piperonyl group, a 3-(1-benzimidazolonyl)propyl group; a group of formula

$$-(CH_2)_{\Pi_{111}}-X^{III}-$$

in which n_{III} is 0, 1, 2 or 3, X^{III} is a single bond or alternatively -O-, -S-, -NH-, -CO-, -CH=CH- or

10

5

and R₃^{III} is H, CH₃, halogen, CN, CF₃ or an acyl group -COR₄^{III}, R₄^{III} being a linear or branched alkyl group having 1 to 6 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms or a phenyl group which can bear a CH₃ or F substituent; or alternatively a group of formula

15

20

in which Z^{III} denotes an O or S atom or a divalent group NH, N-CH₃ or N-CN and R₅^{III} denotes a linear or branched alkyl group having 1 to 8 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms which can bear a phenyl substituent, a (C3-C6 cycloalkyl) (linear or branched, C1-C3 alkyl) group, a phenyl group which can bear a CH₃, halogen or CF₃ substituent, a phenyl(linear or branched, C₁-C₃ alkyl) group or a naphthyl, adamantyl or ptoluenesulphonyl group.

25

[0106] Preferred compounds (III) are those with RIII representing the group

30

 Z^{III} and R^{III}_5 being as above-defined and Z^{III} is especially O, S or NH. [0107] Preferred group R^{III}_5 is a (C₃-C₆)cycloalkyl group. [0108] Preferred R¹ and R² groups are as above-described in formula (A).

[0109] An example of such compound (III) is N'-Cyclohexylthiocarbamoyl-N-1,4'-bipiperidine (compound 84).

101101 According to a fourth aspect, a sub-class of compounds (A) includes the compounds which have the following formula (IV), analogous to compounds disclosed in EP 494 010: in which

40

$$\mathbb{R}^{N} - \mathbb{N} \longrightarrow \mathbb{R}^{1} \qquad (N)$$

45

- R¹ and R² are as defined with reference to general formula (A);
 R^{IV} represents a hydrogen atom or a group COR₃^{IV}, in which R₃^{IV} represents
 - (a) a linear or branched aliphatic group containing 1 to 11, and in particular 1 to 9, carbon atoms;
 - (b) a cyclane ring-system such as cyclopropane, phenylcyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, norbomane, adamantane, noradamantane, chlorooxonorbomane, chloroethylenedioxynorbomane, bromoethylenedioxynorbomane and the anhydride group of hydroxycarboxy-1,2,2trimethylcyclopentanecarboxylic acid;
 - (c) a benzene ring, unsubstituted or substituted at the para-position with a linear or branched aliphatic group containing 3 to 5 carbon atoms, as well as with a halogen;

(d) a group $(CH_2)_{m|V}R_4^{IV}$ in which m_{IV} is a number between 1 and 10, and R_4^{IV} represents a cyclane ring system such as cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexane, cycloheptane, norbomane, noradamantane, adamantane and 6,6-dimethylbicyclo[3.1.1] heptene; a benzene ring, unsubstituted or monosubstituted with a fluorine atom, a chlorine atom, a methyl group or a methoxy group; a thiophene ring grafted via its ring-position 2 or its ring-position 3; a carboxylic acid ester group $COOR_5^{IV}$, in which R_5^{IV} is a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbomane; a carboxylic acid amide group of structure $CONHR_6^{IV}$, in which R_6^{IV} represents a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbornane; a carboxylic acid amide group of structure

CON

in which the group

20

25

30

35

40

55

5

10

15

represents pyrrolidine, piperidine or 2,6-dimethylmorpholine; or an ether group —O-R₇^{IV}, it being possible for R₇^{IV} to be a benzene ring, unsubstituted or monosubstituted with a chlorine or fluorine atom or disubstituted with a chlorine atom and with a methyl group;

(e) a group -CH=CHR₈^{IV}, in which R₈^{IV} represents a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane, norbomane or norbornene;

(f) a secondary amine group -NH(CH₂)_{nIV}R₉^{IV}, in which n_{IV} is a number between 1 and 5 and R₉^{IV} constitutes a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbornane, or a benzene ring, unsubstituted, mono-substituted with a fluorine or chlorine atom or with a methoxy group or trisubstituted with methoxy groups;

RIV also represents a hydroxyalkenyl group

in which p_{IV} is a number between 2 and 9 and R_{10}^{IV} , represents a benzene ring or a phenoxy group; as well as a group

CSNH(CH₃)_{nIV}R₉IV

— in which n_{IV} is a number between 1 and 5 and R₉^{IV} has the meaning stated above.

45 [0111] Preferred compounds (IV) are those in which R^{IV} represents the group COR₃^{IV}, R₃^{IV} representing especially an aliphatic group a).

[0112] An example of compound (IV) is N-Heptanoyl-1,4'-bipiperidine (compound 85).

[0113] According to a fifth aspect, the invention is relative to non-imidazole compounds analogous to those disclosed by Plazzi et al. (Eur. J. Med. Chem. 1995, 30, 881).

50 [0114] Thus, another sub-class of compounds (A) comprises compounds having the following formula (V):

$$\begin{array}{c}
R^{1} \\
N - (CH_{2})_{q\overline{V}} - Z^{V} - S
\end{array}$$
(Va)

OF

 R^1 $N-(CH_2)_{q\overline{V}}$ Z^V (Vb)

in which

5

10

25

30

40

55

- R¹ and R² are as defined with reference to formula (A) in claim 1;

Z^V represents NH, O or S;

15 — X^V represents a hydrogen atom or a lower alkyl

- q_V is 2 to 5.

[0115] Preferred groups R¹ and R² are as above-defined with reference to formula (A).

[0116] Representative example is compound 92.

[0117] According to a sixth aspect, the present invention concerns non-imidazole compounds which are analogous to those disclosed in WO 95/14007.

[0118] Thus, another subclass of compounds (A) includes the compounds having the following formula (VI):

$$R^{1}$$
 R^{2}
 R^{2}
 R^{2}
 R^{2}
 R^{2}
 R^{1}
 R^{2}
 R^{2}
 R^{1}
 R^{2}
 R^{1}
 R^{2}
 R^{1}
 R^{2}
 R^{1}
 R^{2}
 R^{2

wherein:

— A^{VI} is selected from -O-CO-NR $^{1}_{VI}$ -, -O-CO-, -NR $^{1}_{VI}$ -CO-, -NR $^{1}_{VI}$ -, -O-, -NR $^{1}_{VI}$ -, -O-, -CO-NR $^{1}_{VI}$ -, -CO-O-, and -C(=NR $^{1}_{VI}$)-NR $^{1}_{VI}$ -;

— the groups R¹_{VI}, which may be the same or different when there are two or three such groups in the molecule of formula VI, are selected from hydrogen, and lower alkyl, aryl, cycloalkyl, heterocyclic and heterocyclylalkyl groups, and groups of the formula -(CH₂)_{yVI}-G^{VI}, where G^{VI} is selected from CO₂R³_{VI}, COR³_{VI}, COR³_{VI}, COR³_{VI}, OR³_{VI}, SR³_{VI}, NR³_{VI}R⁴_{VI}, heteroaryl and phenyl, which phenyl is optionally substituted by halogen, lower alkoxy or polyhaloloweralkyl, and y_{VI} is an integer from 1 to 3;

— R^2_{VI} is selected from hydrogen and halogen atoms, and alkyl, alkenyl, alkynyl and trifluoromethyl groups, and groups of the formula OR^3_{VI} , SR^3_{VI} and $NR^3_{VI}R^4_{VI}$;

R³_{VI} and R⁴_{VI} are independently selected from hydrogen, and lower alkyl and cycloalkyl groups, or R³_{VI} and R⁴_{VI} together with the intervening nitrogen atom can form a saturated ring containing 4 to 6 carbon atoms that can be substituted with one or two lower alkyl groups;

— the group -(CH₂)_{nVI}-A^{VI}-R¹_{VI} is at the 3- or 4-position, and the group R²_{VI} is at any free position;

- m_{VI} is an integer from 1 to 3;

and n_{VI} is 0 or an integer from 1 to 3.

50 [0119] When used herein, the following terms have the given meanings:

lower alkyl (including the alkyl portions of lower alkoxy) — represents a straight or branched, saturated hydrocarbon chain having from 1 to 6 carbon atoms, preferably from 1 to 4;

lower alkenyl (in R²_{VI}) — represents a straight or branched aliphatic hydrocarbon radical having at least one carbon-to-carbon double bond (preferably in conjugation with the benzene ring that the group R² substitutes) and having from 2 to 6 carbon atoms;

lower alkynyl (in R^2_{VI}) — represents a straight or branched aliphatic hydrocarbon radical having at least one carbon-to-carbon triple bond (preferably in conjugation with the benzene ring that the group R^2 substitutes) and having

from 2 to 6 carbon atoms;

5

10

15

20

25

40

50

55

aryl — represents a carbocyclic group having from 6 to 14 carbon atoms and having at least one benzenoid ring. with all available substitutable aromatic carbon atoms of the carbocyclic group being intended as possible points of attachment, said carbocyclic group being optionally substituted with 1 to 3 Yvi groups, each independently selected from halo, alkyl, hydroxy, loweralkyoxy, phenoxy, amino, loweralkylamino, diloweralkylamino, and polyhaloloweralkyl. Preferred aryl groups include 1-naphthyl, 2-naphthyl and indanyl, and especially phenyl and substituted phe-

cycloalkyl — represents a saturated carbocyclic ring having from 3 to 8 carbon atoms, preferably 5 or 6; halogen — represents fluorine, chlorine, bromine and iodine;

heterocyclic — represents, in addition to the heteroaryl groups defined below, saturated and unsaturated cyclic organic groups having at least one O, S and/or N atom interrupting a carbocyclic ring structure that consists of one ring or two fused rings, wherein each ring is 5-, 6- or 7-membered and may or may not have double bonds that lack delocalized pi electrons, which ring structure has from 2 to 8, preferably from 3 to 6 carbon atoms; e.g., 2- or 3-piperidinyl, 2- or 3-piperazinyl, 2- or 3-morpholinyl, or 2- or 3-thiomorpholinyl;

heteroaryl — represents a cyclic organic group having at least one O, S and/or N atom interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclic group having from 2 to 14, preferably 4 or 5 carbon atoms, e.g., 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-thiazolyl, 2- or

2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, or 3- or 4-pyridazinyl, etc.

[0120] Preferred heteroaryl groups are 2-, 3- and 4-pyridyl;

heterocyclyl-alkyl — represents a heterocyclic group defined above substituting an alkyl group; e.g., 2-(3-piperidinyl)-ethyl, (2-piperazinyl)-methyl, 3-(2-morpholinyl)-propyl, (3-thiomorpholinyl)-methyl, 2-(4-pyridyl)-ethyl, (3-pyridyl)-methyl, or (2-thienyl)-methyl.

[0121] Preferably, AVI is -CH₂-NR¹_{VI}- or especially -C(=NH)-NR¹_{VI}- preferred compounds include those wherein m_{VI} is 1 or 2, and n_{VI} is 0, 1 or 2.

[0122] Other preferred values of A include -O-CO-NR¹_{VI}-, -O-, and -CO-O-. In all these compounds, the groups R¹_{VI} are as defined above, and the side chain is preferably at the 4-position. In compounds of formula VI, one group R¹_{VI} is preferably selected from hydrogen, 2-phenylethyl, 4-chlorophenylmethyl, 4-methoxyphenylmethyl, 4-trifluoromethylphenylmethyl and 4-pyridylmethyl, but is especially 4-chlorophenylmethyl; any other group R¹VI that is present is preferably a hydrogen atom or a methyl group.

[0123] Particularly preferred compounds are those wherein n_{VI} and m_{VI} are each 1, and A^{VI} represents an oxygen atom.

[0124]

 R^1_{VI} is preferably an aryl or -(CH₂)_{yVI}- G^{VI} with G^{VI} being a phenyl. R^1 and R^2 are preferably selected as specified with reference to formula (A). [0125]

Another sub-class of compounds (A) comprises compounds of formula (VI) wherein R¹_{VI} represents an aryl [0126] group, especially a phenyl optionally substituted with a keto substituent, R_{Vl}^2 , n_{Vl} , m_{Vl} and A^{Vl} having the above-meaning.

The keto substituent is as above-defined in YII with reference to compounds (IIa) and (IIb). [0127]

Preferred compounds are those with n_{VI} and m_{VI} being each 1 and A^{VI} being an oxygen atom.

[0129] Examples 88, 89 and 90 are illustrative of compounds VI.

[0130] According to a seventh aspect, the present invention is directed to another sub-class of compounds (A) including non-imidazole compounds having the following formula (VII) which are analogous to compounds disclosed in Clitherow et al. (Bioorg. & Med. Chem. Lett., 6 (7), 833, 1996):

$$R^{1} \qquad (CH_{2})n_{VII} \qquad (CH_{2})_{MVII} \qquad (VII)$$

in which

- R¹ and R² are as defined in reference to formula (A);
- X^{VII}, Y^{VII} and Z^{VII} are identical or different and represent O, N or S;
- --- n_{VII} is varying from 1 to 3;
- m_{VII} is 1 or 2.

5

15

20

25

30

35

40

45

50

55

[0131] n_{VII} is preferably 2 or 3, especially 2 and m_{VI} is preferably 1. [0132] Preferred compounds are those with X^{VII} being 0 and Y^{VII} and Z^{VII} each being N to represent a 1, 2, 4-oxadiazolył group.

[0133] An illustrative compound is given in example 91 and as example 102.

[0134] According to a eighth aspect, the present invention is directed to another sub-class of compounds (A) including the non-imidazole compounds having the following formula (VIII), which are analogous to those disclosed in WO 95/06037:

(VIII)

wherein R^1 and R^2 are as defined with reference to formula (A) and wherein [0135] A^{VIII} is

- 1) a group of the formula $(CH_2)_{mV|II}$, wherein $m_{V|II} = 0.9$; or
- 2) a group of the formula:

wherein R⁵_{VIII} represents hydrogen, (C₁-C₃)alkyl-, aryl(C₁-C₃)alkyl-, aryl-, wherein aryl may optionally be substituted, hydroxyl-, (C_1-C_3) alkoxy-, halogen, amino-, cyano- or nitro; and R^6_{Viii} represents hydrogen, (C_1-C_3) alkyl-, aryl(C1-C3)alkyl-, or aryl-, wherein aryl may optionally be substituted; or

3) a group of the formula:

wherein R5_{VIII} and R6_{VIII} are as defined above; or 4) a group of the formula:

if BVIII is a group of the formula:

-c(

such that A^{VIII} and B^{VIII} together form a group of the formula:

10 R⁶_{VIII}

wherein R⁶_{VIII} is as defined above; or 5) a group of the formula:

5

15

30

35

40

50

20 R⁶vIII C C R⁸vI

wherein R⁶_{VIII} is as defined above; or 6) a group of the formula: if B^{VIII} is a group of the formula:

R⁶√III R⁶√I — Ç — C = R⁶√III

-c(

such that A^{VIII} and B^{VIII} together form a group of the formula:

wherein R^6_{VIII} is as defined above; or 7) a group of the formula:

 $-(CH_2)_{X_{VIII}}-S-(CH_2)_{Y_{VIII}}$

wherein $x_{VIII} + y_{VIII} = m_{VIII}-1$;

B^{VIII} is

1) a group of the formula:

5

10

wherein R^5_{VIII} is as defined above; or 2) a group of the formula:

15

20

if A is a group of one of the formulas:

$$\begin{array}{cccc}
\mathsf{R}^{6}_{\mathsf{VIII}} & & & \mathsf{R}^{6}_{\mathsf{VIII}} \mathsf{R}^{6}_{\mathsf{VIII}} \\
\mathsf{C} & & & & & & & \\
\mathsf{R}^{6}_{\mathsf{VIII}} & & & \\
\mathsf{R}^{6}_{\mathsf{VIII}} & & & &$$

25

such that A and B together form a group of one of the formulas:

30

35

wherein R^6_{VIII} is as defined above; or 3) a group of the formula:

40

if X^{VIII} is a group of the formula:

45

50

such that \mathbf{B}^{VIII} and \mathbf{X}^{VIII} together form a group of the formula

wherein $p_{VIII} = 1-3$; or X^{VIII} is

5

10

15

20

25

30

35

40

45

50

55

1) a group of the formula $(CH_2)_{nVIII}$ wherein $n_{VIII} = 2-4$; or

2) a group of the formula:

if B^{VIII} is a group of the formula:

>c-

such that XVIII and BVIII together form a group of the formula:

wherein $p_{VIII} = 1-3$; or

3) two hydrogens (one on the carbon and one on the nitrogen); or

4) one hydrogen on the carbon atom and one R⁷_{VIII} group on the nitrogen atom,

wherein R⁷_{VIII} represents hydrogen, (C₁-C₁₀)alkyl-, aryl (C₁-C₁₀)alkyl-, or aryl, wherein aryl may optionally be substituted;

 Y^{VIII} is a group of the formula (CH₂)_{kVIII}, wherein k_{VIII} = 0-2;

 R^4_{VIII} represents hydrogen, (C_1-C_{10}) alkyl-, (C_1-C_3) alkyl-sulfonamide-, aryl (C_1-C_{10}) alkyl-, aryl, wherein aryl may optionally be substituted;

or a group of the formula:

or a group of the formula:

wherein XVIII represents O, S, or NH,

R⁷_{VIII} is as defined as above;

R8VIII represents (C1-C10)alkyl-, aryl(C1-C10)alkyl- or aryl,

wherein aryl may optionally be substituted and wherein aryl is phenyl, substituted phenyl, naphtyl, substituted naphtyl, pyridyl.

[0136] The present invention comprises both linear and ringstructured compounds.

101371 The linear compounds have for example one of the formulas

$$R^{1}$$
 $(CH_{2})_{n}^{VIII}$ NH_{2} or R^{2} $(CH_{2})_{n}^{VIII}$ NH C NH R^{VIII} $(VIIIb)$

[0138] Preferred R¹ and R² groups are as defined with reference to formula (A).

[0139] A compound (VIII) is described in example 94.

[0140] According to a ninth aspect, the invention is relative to a sub-class of compounds (A) consisting of compounds having the following formula (IX) which are analogous to those described in WO 97/29092:

$$\begin{array}{c|c}
R^1 & R^2_{ix} \\
R^2 & R^1_{ix}
\end{array}$$
(IX)

wherein:

5

10

20

35

40

45

50

R¹ and R² are as defined with reference to formula (A)

 R^1_{IX} is C_4 to C_{20} hydrocarbyl (in which one or more hydrogen atoms may be replaced by halogen, and up to four carbon atoms [and especially from 0 to 3 carbon atoms] may be replaced by oxygen, nitrogen or sulphur atoms, provided that R^1_{IX} does not contain an -O-O-group),

R²_{IX} is H or C₁ to C₁₅ hydrocarbyl (in which one or more hydrogen atoms may be replaced by halogen, and up to three carbon atoms may be replaced by oxygen, nitrogen or sulphur atoms, provided that R²_{IX} does not contain an -O-O-oroup).

 $m_{|X}$ is from 1 to 15 (preferably 1 to 10, more preferably 3 to 10, eg. 4 to 9) each $X^{|X|}$ group is independently

or one X^{IX} group is -N(R^4_{IX})-, -O- or -S- (provided that this X^{IX} group is not adjacent the -NR $^2_{IX}$ - group) and the remaining X^{IX} groups are independently

wherein

 R_{IX}^3 is H, C_1 to C_6 alkyl, C_2 to C_6 alkenyl, $-CO_2R_{IX}^5$, $-CON(R_{IX}^5)_2$, $-CR_{IX}^5$ OR $_{IX}^6$ or $-OR_{IX}^5$ (in which R_{IX}^5 and R_{IX}^6 are H or C_1 to C_3 alkyl), and R_{IX}^4 is H or C_1 to C_6 alkyl.

[0141] The term "hydrocarbyl", as used herein, refers to monovalent groups consisting of carbon and hydrogen. Hydrocarbyl groups thus include alkyl, alkenyl, and alkynyl groups (in both straight and branched chain forms), cycloalkyl (including polycycloalkyl), cycloalkenyl, and aryl groups, and combinations of the foregoing, such as alkylaryl,

alkenylaryl, alkynylaryl, cycloalkylaryl, and cycloalkenylaryl groups.

[0142] A "carbocyclic" group, as the term is used herein, comprises one or more closed chains or rings, which consist entirely of carbon atoms. Included in such groups are alicyclic groups (such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and adamantyl), groups containing both alkyl and cycloalkyl moieties (such as adamantanemethyl), and aromatic groups (such as phenyl, naphthyl, indanyl, fluorenyl, (1,2,3,4)-tetrahydronaphthyl, indenyl and isoindenyl).

[0143] The term "aryl" is used herein to refer to aromatic carbocyclic groups, including those mentioned above.

[0144] When reference is made herein to a substituted carbocyclic group (such as substituted phenyl), or a substituted heterocyclic group, the substituents are preferably from 1 to 3 in number and selected from C1 to C6 alkyl, C1 to C6 alkoxy, C1 to C6 alkylthio, carboxy, C1 to C6 carboalkoxy, nitro, trihalomethyl, hydroxy, amino, C1 to C6 alkylamino, di(C₁ to C₆ alkyl)amino, aryl, C₁ to C₆ alkylaryl, halo, sulphamoyl and cyano.

[0145] The term "halogen", as used herein, refers to any of fluorine, chlorine, bromine and iodine.

[0146] Preferably, R^2_{IX} is selected from H, C_1 to C_6 alkyl, C_1 to C_6 cycloalkyl, C_1 to C_6 hydroxyalkyl, C_1 to C_6 alkyl and substituted aryl C_1 to C_6 alkyl. For example, R^2_{IX} may be H or C_1 to C_3 alkyl.

[0147] In certain embodiments, -X^{IX}_{mIX} is a C₁ to C₈ alkylene group, e.g. a butylene group.
[0148] Included in the definition of R¹_{IX} are aryl-containing groups (such as phenyl, substituted phenyl, naphthyl and substituted naphthyl), and (cycloalkyl)alkyl groups (such as cyclohexylpropyl and adamantylpropyl).

[0149] Preferably, R¹_{IX} is a group of the formula

$$R_{i}^{11}_{ix}$$
 $R_{i}^{13}_{ix}$ $R_{i}^{12}_{ix}$ $R_{i}^{12}_{ix}$

wherein 25

20

piX is 0 or 1,

 R^{11}_{IX} is H or C_1 to C_3 alkyl,

q_{IX} is from 0 to 4,

R¹²IX is a carboxyclic, substituted carbocyclic, heterocyclic or substituted heterocyclic group, and

 R^{13}_{IX} is independently selected from H, C_1 to C_6 alkyl, C_1 to C_6 cycloalkyl, C_1 to C_6 hydroxyalkyl, C_1 to C_6 alkylhydroxyalkyl, aryl C_1 to C_6 alkyl and substituted aryl C_1 to C_6 alkyl.

[0150] Preferably, R¹³_{IX} is hydrogen.

[0151] Compounds (IX) wherein R¹_{IX} is a group -NH-CH₂-Ph where Ph represents an optionally substituted phenyl, are preferred.

[0152] Preferred groups R¹ and R² are as specified with reference to formula (A).


[0153] An illustrative example is compound 100.

[0154] According to a tenth aspect, the present invention is relative to another sub-class of compounds (A) comprising compounds having the following formula (X), which are analogous to compounds disclosed by Wolin et al. (Bioorg. & Med. Chem. Lett., 8, 2157 (1998)):

40

45

50



wherein:

- R¹ and R² are as defined with reference to formula (A);
- R¹_x is H or CH₃;
- R²_x is selected from a phenyl optionally substituted with a halogen atom, preferably chlorine, a (C₁-C₄)alkyl, a (C₁-C₄)alkoxy, CF₃, OCF₃, NO₂, NH₂; or a CH₂-phenyl optionally substituted as above-specified;
- n_x is from 0 to 3.

[0155] n_x is preferably 1. R² is preferably a phenyl group, especially a mono-substituted phenyl group.

[0156] Preferred R¹ and R² are as above-specified for formula (A).

[0157] Compound 101 is illustrative of compounds (X).

[0158] According to a eleventh aspect, the invention is directed to non-imidazole compounds which are analogous to those disclosed in WO 96/38142.

[0159] Thus, another sub-class of compounds (A) of the invention is directed to compounds having the following formula (XI):

$$\begin{array}{c|c}
R^{1} & X^{XI} \\
R^{2} & X^{XI} \\
R_{2}^{XI} & (CH_{2})_{\Pi} X^{I} \\
R_{1}^{XI} & (XI)
\end{array}$$

where R¹ and R² are as defined with reference to formula (A);

where A^{XI} is -NHCO-, -N(CH₃)-CO-, -NHCH₂-, -N(CH₃)-CH₂-, -CH=CH-, -COCH₂-, CH₂CH₂-, -CH(OH)CH₂-, or -C=C-; X^{XI} is H, CH₃, NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, or SH;

R₂^{XI} is hydrogen or a methyl or ethyl group;

R₃XI is hydrogen or a methyl or ethyl group;

nXI is 0, 1, 2, 3, 4, 5 or 6; and

10

R₁^{XI} is selected from the group consisting of C₃ to C₈ cycloalkyl; phenyl or substituted phenyl; decahydronaphthalene and octahydroindene; or

R₁XI and XXI may be taken together to denote a 5,6 or 6,6 saturated bicyclic ring structure when XXI is NH, O, S, or SO₂. [0160] Preferably for compounds of formula (XI):

AXI is -NHCO-, -N(CH₃)-CO-, -NHCH₂-, -N(CH₃)-CH₂-, -CH=CH-, -COCH₂-, -CH₂CH₂-, -CH(OH)CH₂-, or -C=C-;

 X^{XI} is H, CH₃, NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, or SH;

R₂XI is hydrogen or a methyl or ethyl group;

R₃^{XI} is hydrogen or a methyl or ethyl gorup;

n^{XI} is 0, 1, 2, 3, 4, 5, or 6; and

R₁XI is selected from the group consisting of (a) C₃ to C₈ cycloalkyl; (b) phenyl or substituted phenyl; (d) heterocyclic (e) decahydronaphthalene and (f) octahydroindene; or

R₁XI and XXI may be taken together to denote a 5,6 or 6,6 saturated bicyclic ring structure when XXI can be NH, O, or S. [0161] More preferably, the present invention provides compounds

where AXI is -NHCH2-, -N(CH3)-CH2-, -CH=CH-,

-COCH₂-, -CH₂CH₂, -CH(OH)CH₂-, or -C=C-;

XXI is H, CH3, NH2, NH(CH3), N(CH3)2, OH, OCH3, or SH;

RXI2 is hydrogen or a methyl or ethyl group;

RXI3 is hydrogen or a methyl or ethyl group;

n^{XI} is 0, 1, 2, 3, 4, 5, or 6; and

RXI₁ is selected from the group consisting of (a) C₃ to C₈ cycloalkyl; (b) phenyl or substituted phenyl; (d) heterocyclic; (e) decahydronaphthalene and (f) octahydroindene; or

RXI and XXI may be taken together to denote a 5,6 or 6,6 saturated bicyclic ring structure when XXI can be NH, O, or S. [0162] Most preferably, the present invention provides compounds

where AXI is -CH=CH or -C=C-;

 X^{XI} is H, CH_3 or NH_2 ; R_2^{XI} and R_3^{XI} are H;

n^{Xi} is 1, 2, or 3;

R₁XI is selected from the group consisting of (a) C₃ to C₈ cycloalkyI; (b) phenyl or substituted phenyl; (d) heterocyclic; (e) decahydronaphthalene and (f) octahydroindene; or

 R_1^{XI} and X^{XI} may be taken together to denote a 5,6 or 6,6 saturated bicyclic ring structure when X^{XI} is NH, O, or S.

[0163] The term "substituted phenyl" as used herein refers to a phenyl group substituted by one or more groups such as alkyl, halogen, amino, methoxy and cyano groups.

[0164] The term "alkyl" refers to straight or branched chain radicals. Representative examples of alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl and the like.

[0165] Compounds (XI) where AXI is -CH=CH- or —C=C-, XXI, R2XI and R3XI are each H, nXI is 1 and R1XI is a C3-C_B cycloalkyl, are especially preferred.

[0166] R¹ and R² are preferably selected as above-indicated in reference to formula (A).

[0167] Representative particularly preferred compounds are compounds 104, 105 or 106.

[0168] According to a twelfth aspect, the invention concerns non-imidazole compounds which are analogous to those disclosed in WO 96/38141.

[0169] Thus, the invention is relative to compounds having the following formula (XII):

(XII)

where R1 and R2 are as defined in reference to formula (A),

where R_2^{XII} is a hydrogen or a methyl or ethyl group; R_3^{XII} is a hydrogen or a methyl or ethyl group;

n^{XII} is 0, 1, 2, 3, 4, 5, or 6; and

 R_1^{XII} is selected from the group consisting of (a) C_3 to C_8 cycloalkyl; (b) phenyl or substituted phenyl; (c) alkyl; (d) heterocyclic; (e) decahydronaphthalene; and (f) octahydroindene; with the provisos that

when X^{XII} is H, A^{XII} can be -CH₂CH₂-, -COCH₂-, -CONH-, -CON(CH₃)-, -CH=CH-, -C≡C-, -CH₂-NH-, -CH₂-N(CH₃)-, -CH(OH)CH₂-, -NH-CH₂-, -N(CH₃)-CH₂-, -CH₂O-, -CH₂S-, or —NHCOO-; when XXII is NH2, NH(CH3), N(CH3)2, OH, OCH3, CH3, SH or SCH3; AXII can be -NHCO-, -N(CH3)-CO-, -NHCH2-, -N(CH₃)-CH₂-, -CH=CH-, -COCH₂-, -CH₂CH₂-, -CH(OH)CH₂-, -Cr-Cr-; and when R₁XIII and XXIII taken together denote a 5,6 or 6,6 saturated bicyclic ring structure XXIII can be NH, O, or S.

[0170] The term "alkyl" as used herein refers to straight or branched chain radicals derived from saturated hydrocarbons by the removal of one hydrogen atom. Representative examples of alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, and the like.

[0171] The term "substituted phenyl" as used herein refers to a phenyl group substituted by one or more groups such as alkyl, halogen, amino, methoxy, and cyano groups.

[0172] The term "bicyclic alkyl" as used herein refers to an organic compound having two ring structures connected to an alkyl group. They may or may not be the same type of ring and the rings may be substituted by one or more groups. Representative bicyclic alkyl groups include adamanthyl, decahydronaphthalene and norbornane.

[0173] The cyclopropane attached to the NR¹R² moiety is preferably in trans configuration.

[0174] More preferably, the present invention provides compounds of the general formula (XII):

where AXII is -CONH, -CH=CH-, -NHCOO-, or -C=C-;

XXII is H or NH2;

 R_2^{XII} and R_3^{XII} are H; n^{XII} is 0, 1, 2 or 3;

R₁^{XII} is cyclohexyl, phenyl or substituted phenyl.

[0175] In compounds (XII), AXII is especially -CH=CH- or -C=C-;

R₂XII, R₃XIII and XXIII are each especially a hydrogen atom;

n_{XII} is preferably 1 and R₁^{XII} is especially an alkyl group.

[0176] R¹ and R² are preferably selected as above-indicated with reference to formula (A).

[0177] Representative example of compounds (XII) is compound 107.

[0178] According to a thirteenth aspect, the invention is directed to non-imidazole compounds analogous to those disclosed in WO 95/11894.

[0179] Thus, the present invention is relative to a sub-class of compounds (A) comprising compounds having the following formula (XIII):

-50

5

$$R^{1} \longrightarrow D^{\times |||} (O) x_{X|||} (CH_{2})_{DX|||} R_{2}^{X|||}$$

$$(X||||)$$

5

10

wherein R^1 and R^2 are as defined with reference to formula (A) wherein D^{XIII} is CH_2 or CH_2 - CH_2 , Z^{XIII} represents sulfur (S) or oxygen (O), preferably O, X_{XIII} is 0 or 1, n_{XIII} is an integer

and R2XIII represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms including mono and bicyclic moieties, and a substituted or an unsubstituted aryl group of up to about 20 carbon atoms, or any combination of abovementioned groups, or salts thereof.

[0180] In a specific embodiment, R₂XIII can represents a disubstituted methyl, such as but not limited to dicyclohexyl methyl (-CH(C_6H_{11})₂), diphenyl methyl (-CH(C_6H_5)₂), and the like. If R_2^{XIII} is tert-butyl, cyclohexyl, or dicyclohexylmethyl, X_{XIII} or n_{XIII} must not be 0. If R_2^{XIII} is adamantane, the sum of x_{XIII} and n_{XIII} must be greater than 1.

[0181] In a preferred embodiment, DXIII is CH2-CH2, resulting in a piperidine ring structure. However, it is contemplated that DXIII can be CH2, yielding a pyrrolidine ring structure. In yet another embodiment, DXIII can be (CH2)3, yielding a cycloheptimide (seven membered heterocycle with one nitrogen).

[0182] In a specific embodiment, a tetramethylene bound to the amide or carbamate group is used. Preferably a cyclic alkyl or aryl group is linked to the amide or carbamate via the straight chain alkyl group. In a specific embodiment, tetramethylene cyclohexane (cyclohexylbutyl) is bound to an amide. Although specific hydrophobic alkyl and aryl groups have been mentioned, one of ordinary skill in the art will recognize that there are many possible hydrophobic groups for use in the compounds of the invention. These fall within the scope of the instant invention.

[0183] Thus, R₂XIII can be one or more bulky substituent groups. As stated above, in a preferred aspect of the invention, the bulky substituents are removed from the amide or carbamate group on the piperidyl, by increasing $n_{X|II}$. In one embodiment, R_2^{XIII} is $CHR_3^{XIII}R_4^{XIII}$, in which $n_{X|II}$ is 3 or 4 and R_3^{XIII} and R_4^{XIII} are cyclohexyl, phenyl, or the like. R_3^{XIII} and R_4^{XIII} can be the same group or different groups. In another embodiment, R_2^{XIII} is decalin or adamantane or the like. If R_2^{XIII} is adamantane, preferably n_{XIII} is greater than 1, but the sum of x_{XIII} and n_{XIII} must be greater than 1.

[0184] As used herein, the phrase linear chain or branched chained alkyl groups of up to about 20 carbon atoms means any substituted or unsubstituted acyclic carbon-containing compounds, including alkanes, alkenes and alkynes. Examples of alkyl groups include lower alkyl, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tertbutyl; upper alkyl, for example, octyl, nonyl, decyl, and the like; and lower alkylene, for example, ethylene, propylene, propyldiene, butylene, butyldiene, and the like. The ordinary skilled artisan is familiar with numerous linear and branched alkyl groups, which are with the scope of the present invention.

[0185] In addition, such alkyl group may also contain various substituents in which one or more hydrogen atoms has been replaced by a functional group. Functional groups include but are not limited to hydroxyl, amino, carboxyl, amide, ester, ether, and halogen (fluorine, chlorine, bromine and iodine), to mention but a few.

[0186] As used herein, substituted and unsubstituted carbocyclic groups of up to about 20 carbon atoms means cyclic carbon-containing compounds, including but not limited to cyclopentyl, cyclohexyl, cycloheptyl, admantyl, and the like. Such cyclic groups may also contain various substituents in which one or more hydrogen atoms has been replaced by a functional group. Such functional groups include those described above, and lower alkyl groups as describe above. The cyclic groups of the invention may further comprise a heteroatom. For example, in a specific embodiment, R₂XIII is

[0187] As used herein, substituted and unsubstituted aryl groups means a hydrocarbon ring bearing a system of conjugated double bonds, usually comprising six or more even number of $\pi(pi)$ electons. Examples of aryl groups include, by are not limited to, phenyl, naphthyl, anisyl, toluyl, xylenyl and the like. According to the present invention, aryl also includes heteroaryl groupss, e.g., pyrimidine or thiophene. These aryl groups may also be substituted with any number of a variety of functional groups. In addition to the functional groups described above in connection with substituted alkyl groups and carbocyclic groups, functional groups on the aryl groups can be nitro groups.

[0188] As mentioned above, R₂XIII can also represents any combination of alkyl, carbocyclic or aryl groups, for example, 1-cyclohexylpropyl, benzyl cyclohexylmethyl, 2-cyclohexylpropyl, 2,2-methylcyclohexylpropyl, 2,2-methylphenylpropyl, 2,2-methylphenylbutyl.

In a specific embodiment, R2 represents cyclohexane, and nxIII=4 (cyclohexylvaleroyl). In another specific embodiment, R2XIII represents cinnamoyl.

[0190] Particularly preferred are compounds (XIII) wherein ZXIII is an oxygen atom and wherein xXIII is 0 or 1, nXIII is an integer from 0 to 6, more preferably $n_{XIII} = 3-6$, and most preferably $n_{XIII} = 4$, and R_2^{XIII} is as defined above. Examples of preferred alkyl groups for R₂XIII include but are not limited to cyclopentyl, cyclohexyl, admantane methylene, dicyclohexyl methyl, decanyl and t-butyryl and the like. Examples of preferred aryl and substituted aryl groups include but are not limited to phenyl, awl cyclohexyl methyl and the like.

[0191] Preferred R¹ and R² are selected as indicated with reference to formula (A).

[0192] Representative example is compound 103.

[0193] According to a fourteenth aspect, the present invention is directed to compounds analogous to those disclosed in WO 93/12107.

[0194] Thus, a sub-class of compounds (A) of the invention concerns compounds having the following formula (XIV)

$$\begin{array}{c|c}
R^{1}_{XIV} & R^{2}_{XIV} \\
R^{1}_{XIV} & C)_{m^{XIV}} & C)_{m^{XIV}} & R^{3}_{XIV} \\
R^{2}_{XIV} & C)_{m^{XIV}} & C^{2}_{XIV} & C^{2}_{XIV} \\
R^{2}_{XIV} & C^{2}_{XIV} & C^{2}_{XIV} & C^{2}_{XIV}
\end{array}$$
(XIV)

wherein R1 and R2 are as defined in reference of formula (A);

(A) m_{XIV} is an integer selected from the group consisting of: 1 and 2;

(B) n_{XIV} and p_{XIV} are intergers and are each independently selected from the group consisting of: 0, 1, 2, 3, and 4 such that the sum of n_{XIV} and p_{XIV} is 4 and T^{XIV} is a 6-membered ring;

- (C) R^3_{XIV} and R^4_{XIV} are each independently bound to the same or different carbon atom of ring T^{XIV} such that there is only one R^3_{XIV} group and one R^4_{XIV} group in ring T^{XIV} , and each R^1_{XIV} R^2_{XIV} and R^4_{XIV} is independently selected from the group consisting of:

15

20

25

30

35

40

45

- (2) C₁ to C₆ akyl; and
- (3) -(CH₂)_{qXIV}-R⁶_{XIV} wherein q_{XIV} is an integer of: 1 to 7, and R⁶_{XIV} is selected from the group consisting of: phenyl, substituted phenyl, -OR⁷_{XIV}, -C(O)OR⁷_{XIV}, -C(O)OR⁷_{XIV}, -C(O)NR⁷_{XIV}, -C(O)NR⁷_{XIV}, CN and -SR7_{XIV} wherein R7_{XIV} and R8_{XIV} are as defined below, and wherein the substituents on said substituted phenyl are each independently selected from the group consisting of: -OH, -O-(C₁ to C₆)alkyl, halogen, C₁ to C₆ alkyl, -CF₃, -CN, and -NO₂, and wherein said substituted phenyl contains from 1 to 3 substituents;
- (D) R⁵_{XIV} is selected from the group consisting of:

 - (2) C₁ to C₂₀ alkyl;
 - (3) C₃ to C₆ cycloalkyl;
 - (4) -C(O)OR⁷⁷XIV; wherein R⁷⁷XIV is the same as R⁷XIV defined below except that R⁷¹XIV is not H;

 - (5) -C(O)R^{7'}_{XIV}; (6) -C(O)NR^{7'}_{XIV}R⁸_{XIV};
 - (7) allyl;

 - (9) -(CH₂)_q-R⁶_{XIV} wherein q_{XIV} and R⁶_{XIV} are as defined above, and when q_{XIV} is equal to 1, then R⁶_{XIV} is not
- (E) R^7_{XIV} and R^8_{XIV} are each independently selected from the group consisting of: H, C_1 to C_6 alkyl, and C_3 to C_6
- (F) the dotted line (-----) represents a double bond that is optionally present when mxIV is 1, and nXIV is not 0, and p is not 0 (i.e., the nitrogen in the ring is not bound directly to the carbon atom bearing the double bond), and when said double bond is present then R2XIV is absent; and

(G) when m_{XIV} is 2, each R¹_{XIV} is the same or different substituent for each m_{XIV} and each R²_{XIV} is the same or different substituent for each m_{XIV}, and at least two of the substituents R¹_{XIV} and/or R²_{XIV} are H.

[0195] Those skilled in the art will appreciate that the total number of substituents on each of the -(C)_nXIV₋ and -(C) groups is two, and that such substituents are independently selected from the group consisting of hydrogen, R^3_{XIV} and R^4_{XIV} such that there is a total of only one R^3_{XIV} and one R^4_{XIV} substituent in ring T^{XIV} .

[0196] As used herein the following terms have the following meanings unless indicated otherwise:

alkyl — represents a straight or branched, saturated hydrocarbon chain having from 1 to 20 carbon atoms; cycloalkyl — represents a saturated carbocyclic ring having from 3 to 6 carbon atoms; halogen (halo) — represents fluoro, chloro, bromo or iodo.

[0197] Preferably, for compounds of formula (XIV) m is 1; R^5_{XIV} is selected from the group consisting of H and C_1 to C_{15} alkyl; and R^1_{XIV} to R^4_{XIV} are each independently selected from the group consisting of: H, C_1 to C_6 alkyl, and - $(CH_2)_{qXIV}$ - R^6_{XIV} wherein R^6_{XIV} is phenyl. Most preferably, R^5_{XIV} is selected from the group consisting of H and C_1 to C_6 alkyl with H and methyl being even more preferable; and R^3_{XIV} and R^4_{XIV} are each independently selected from the group consisting of: H and methyl.

[0198] Representative compounds of this invention include compounds of the formula:

For formula (XIVa), (XIVb) or (XIVc), R⁵_{XIV} is preferably H or CH₃; R³_{XIV} and R⁴_{XIV} are preferably each an hydrogen atom.

[0200] Preferred R¹ and R² are as specified for formula (A).

[0201] According to a fifteenth aspect, the invention is directed to compounds analogous to those disclosed in WO 93/12108.

[0202] Thus, the invention concerns compounds having the following formula (XV):

55

45

$$R^{1}_{XV}$$
 R^{2}_{XV} R^{3}_{XV} R^{4}_{XV} R^{1}_{XV} R^{2}_{XV} R^{2}_{XV} R^{3}_{XV} R^{4}_{XV} R^{5}_{XV} R^{5}_{XV} R^{6}_{XV} R^{6}_{XV}

wherein R1 and R2 are as defined in reference to formula (A)

- (A) m_{XV} is an integer selected from the group consisting of: 0,1, and 2;
- (B) n_{XV} and p_{XV} are integers and are each independently selected from the group consisting of: 0, 1, 2, and 3 such that the sum of n_{XV} and p_{XV} is 2 or 3 such that when the sum of n_{XV} and p_{XV} is 2, T^{XV} is a 4-membered ring and when the sum of n_{XV} and p_{XV} is 3, T^{XV} is a 5-membered ring;
- (C) each R¹_{XV} R²_{XV} R³_{XV} R⁴_{XV} R⁶_{XV} R⁷_{XV} and R⁸_{XV} is independently selected from the group consisting of:
 - (1) H;

5

10

15

20

25

30

35

40

50

- (2) C₁ to C₆ alkyl;
- (3) C₃ to C₆ cycloalkyl; and
- (4) -(CH₂) $_{q}^{XV}$ -R⁹ $_{XV}$ wherein q $_{XV}$ is an integer of: 1 to 7, and R⁹ $_{XV}$ is selected from the group consisting of: phenyl, substituted phenyl, -OR¹⁰ $_{XV}$ -C(O)OR¹⁰ $_{XV}$ -C(O)R¹⁰ $_{XV}$ -C(O)NR¹⁰ $_$ nyl are each independently selected from the group consisting of: -OH, -O-(C1 to C6) alkyl, halogen, C1 to C6 alkyl, -CF3, -CN, and -NO2, and wherein said substituted phenyl contains from 1 to 3 substituents; examples of -(CH₂)_{qXV}-R⁹_{XV} include benzyl, substituted benzyl and the like, wherein the substitutents on the substituted benzyl are as defined above for said substituted phenyl;
- (D) R⁵_{XV} is selected from the group consisting of:
 - (1) H;
 - (2) C₁ to C₂₀ alkyl;
 - (3) C₃ to C₆ cycloalkyl;
 - (4) -C(O)OR $^{10'}$ XV; wherein R $^{10'}$ XV is the same as R 10 XV defined below except that R $^{10'}$ XV is not H;

 - (5) -C(O)R¹⁰XV; (6) -C(O)NR¹⁰XVR¹¹XV;
 - (7) allyl;

 - (9) -(CH₂) $_{q}^{XV}$ -R⁹ $_{XV}$ wherein q_{XV} and R⁹ $_{XV}$ are as defined above with the proviso that when q_{XV} is 1 then R⁹ $_{XV}$ is not -OH or -SH:
- (E) R¹⁰_{XV} and R¹¹_{XV} are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and C₃ to C₆ cycloalkyl; and, for the substituent -C(O)NR¹⁰XVRXV¹¹, R¹⁰XV and R¹¹XV together with the nitrogen to which they 45 are bound, can form a ring having 5, 6, or 7 atoms;
 - (F) the dotted line (----) represents a double bond that is optionally present when m_{XV} is 1, and T^{XV} is a 5-membered ring, and n_{XV} is not 0, and p_{XV} is not 0 (i.e., the nitrogen in the ring is not bound directly to the carbon atom bearing the double bond), and when said double bond is present then R²XV and R⁸XV are absent;
 - (G) when m_{XV} is 2, each R^1_{XV} is the same or different substituent for each m_{XV} , and each R^2_{XV} is the same or different substituent for each mXV;
 - (H) when n_{XV} is 2 or 3, each R^3_{XV} is the same or different substituent for each n_{XV} and each R^4_{XV} is the same or different substituent for each nXV; and
 - (I) when p_{XV} is 2 or 3, each R^6_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substitutent for each p, and each R^7_{XV} is the same or different substitutent for each p, and each R^7_{XV} is the same or different substitutent for each R^7_{XV} is the same of $R^7_{$ ferent substituent for each pxv
 - [0203] As used herein the following terms have the following meanings unless indicated otherwise:

alkyl — represents a straight or branched, saturated hydrocarbon chain having from 1 to 20 carbon atoms; cycloalkyl — represents a saturated carbocyclic ring having from 3 to 6 carbon atoms; and halogen (halo) —represents fluoro, chloro, bromo or iodo.

- [0204] Preferably, for compounds of formula (XV) m_{XV} is 0 or 1; R⁵_{XV} is selected from the group consisting of H and C₁ to C₂₀ alkyl; and R¹_{XV} to R⁴_{XV} and R⁶_{XV} to R⁸_{XV} are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and -(CH₂)_{qXV} R⁹_{XV} wherein R⁹_{XV} is phenyl. Most preferably, R⁵_{XV} is selected from the group consisting of H and methyl; and R¹_{XV} R²_{XV} R³_{XV} R⁴_{XV} R⁶_{XV} R⁷_{XV} and R⁸_{XV} are each independently selected from the group consisting of: H, methyl, ethyl, pentyl, benzyl, and 2-phenylethyl.
- 10 [0205] Representative compounds of this invention include compounds of the formula:

15

20

25

30

35

45

$$R^{1}_{XV}$$
 R^{2}_{XV}
 R^{2}_{XV}
 R^{3}_{XV}
 R^{4}_{XV}
 R^{5}_{XV}
 R^{5}_{XV}
 R^{6}_{XV}
 R^{6}_{XV}
 R^{6}_{XV}
 R^{6}_{XV}

$$\begin{array}{c|c}
R^{1}_{XV} & R^{2}_{XV} & R^{6}_{XV} \\
R^{1} & R^{2}_{XV} & R^{8}_{XV} & R^{8}_{XV}
\end{array}$$

$$\begin{array}{c|c}
R^{1}_{XV} & R^{6}_{XV} & (XVb) \\
R^{7}_{XV} & R^{8}_{XV} & (XVb)
\end{array}$$

wherein m_{XV} and R^1_{XV} to R^8_{XV} are as defined for formula (XV)

[0206] Compounds (XVc) or (XVd) are preferred.

Representative compounds (XVa) to (XVd) are those wherein R^5_{XV} is H or CH₃. Preferably, only one or two of substituents R^3_{XV} , R^4_{XV} , R^6_{XV} , R^7_{XV} , R^8_{XV} is different from H and represents

[0209] R¹ and R² are preferably selected as indicated in reference to formula (A).

[0210] According to a sixteenth aspect, the invention is directed to compounds analogous to those disclosed in WO

[0211] Thus, the invention is relative to a sub-class of compounds (A) consisting of compounds having the following formula (XVI)

> R1 ZXVI R2XVIR1XVI (IVX)

wherein R¹ and R² are as defined in reference to formula (A)

 Z^{XVI} is a group of the formula (CH₂)_{mXVI} wherein $m_{XVI} = 1-5$ or a group of the formula:

wherein

10

15

20

25

 $R^6_{XVI} = (C_1 - C_3)alkyl$

 $R^{7}_{XVI} = (C_{1}-C_{3})$ alkyl; wherein Z^{XVI} may optionally comprise other substituents selected such that the activity of the derivative is not negatively

XXVI represents S, NH or CH2

 R^1_{XVI} represents hydrogen, (C₁-C₃)alkyl-, aryl(C₁-C₁₀)alkyl, wherein aryl may optionally be substituted, aryl, (C₅-C₇)cycloalkyl(C₁-C₁₀)alkyl-, or a group of the formula:

wherein $n_{XVI} = 1-4$, R^8_{XVI} is aryl, aryl(C_1-C_{10})alkyl-, (C_5-C_7)cycloalkyl- or (C_5-C_7) cycloalkyl(C_1-C_{10})alkyl-, and R^9_{XVI} is hydrogen, (C_1-C_{10})alkyl- or aryl; R_2^{XVI} and R_5^{XVI} represent hydrogen, (C_1-C_3)alkyl-, aryl or arylalkyl-, wherein aryl may optionally be substituted; wherein aryl is phenyl, substituted phenyl, naphthyl, substituted napththyl, pyridyl or substituted pyridyl;

R₂^{XVI} and R₅^{XVI} are preferably a hydrogen atom.

m_{XVI} is preferably 2 or 3 X^{XVI} is preferably S or NH

R₁^{XVI} is preferably selected from H or an optionally substituted aryl.

[0212] Preferred R¹ and R² are selected as specified for formula A.

[0213] According to a seventeenth aspect, a sub-class of compounds (A) of the invention comprises compounds having the following formula (XVII), which can be considered as analogous to those disclosed in EP 680 960:

$$\mathbb{R}^{2}$$
 \mathbb{R}^{4} \mathbb{R}^{4}

[0214] The lower alkyl groups are preferably linear or branched alkyl groups having 1 to 6 carbon atoms. Specific examples thereof include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl and n-hexyl groups.

[0215] The linear or branched alkyl groups are preferably those having 1 to 8 carbon atoms. Specific examples thereof include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl and 1,2,2-trimethylpropyl groups.

[0216] The cycloalkyl groups are preferably those having 3 to 10 carbon atoms. The cycloalkyl groups include not only monocycloalkyl groups (for example, cyclopentyl, cyclohexyl and cycloheptyl) but also polycycloalkyl groups (for example, bicycloalkyl and tricycloalkyl). Examples of the bicycloalkyl groups include norbornyl (for example, exo-2-norbornyl and endo-2-norbornyl), 3-pinanyl and bicyclo[2.2.2]oct-2-yl groups, while examples of the tricycloalkyl groups include adamantyl groups (for example, 1-adamantyl and 2-adamantyl). Such a cycloalkyl group may be substituted by alkyl group(s), etc.

[0217] The cycloalkylalkyl groups are preferably those composed of a cycloalkyl group having 3 to 10 carbon atoms with a linear or branched alkyl group having 1 to 3 carbon atoms. Specific examples thereof include 1-cyclohexylethyl and 1-cyclopropylethyl groups.

[0218] Wherein m_{XVII} represents an integer of from 4 to 6.

5

10

[0219] R 4 _{XVII} represents a hydrogen atom, a linear or branched alkyl group, a cycloalkyl group, and Z^{XVII} represents R 5 _{XVII} or A^{XVII}-R 6 _{XVII}, wherin A^{XVII} represents S or O, R $_5$ ^{XVII} represents a hydrogen atom, a lower alkyl group, a substituted or unsubstituted aralkyl group, and R $_6$ ^{XVII} represents a lower alkyl group, a lower alkyl group;

[0220] The lower alkenyl groups are preferably linear or branched alkenyl groups having 3 to 6 carbon atoms. Specific examples thereof include allyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, cis-2-butenyl, trans-2-butenyl and 3-methyl-2-butenyl groups.

[0221] The lower alkynyl groups are preferably those having 3 to 6 carbon atoms. A specific example thereof includes a 2-propynyl group.

[0222] The substituted aryl groups are preferably phenyl and naphthyl groups which may be substituted by halogen atoms and trifluoromethyl, lower alkyl, lower alkoxy, lower alkylthio, cyano and nitro groups.

[0223] Specific examples thereof include phenyl, 1-naphthyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 4-fluorophenyl, 4-fluorophenyl, 2-methoxyphenyl, 4-methoxyphenyl, 2-tolyl and 3-tolyl groups.

[0224] The aralkyl groups are preferably benzyl and trityl groups.

[0225] The substituted aralkyl groups are preferably arylalkyl groups composed of a phenyl or naphthyl group, which may be substituted by halogen atoms and trifluoromethyl, lower alkyl, lower alkoxy, lower alkylthio, cyano and nitro groups, and a linear or branched alkyl group having 1 to 4 carbon atoms.

[0226] Specific examples thereof include benzyl, α-methylbenzyl, phenethyl, 3-phenylpropyl, 4-phenylbutyl, 4-chlorobenzyl, 4-fluorobenzyl, 4-methoxybenzyl, 4-chloro-α-methylbenzyl, 4-fluoro-αmethylbenzyl and 4-methoxy-α-methylbenzyl groups.

[0227] Among the compounds represented by the general formula (XVII) preferable examples include those wherein: m_{XVII} is from 4 to 6;

R⁴_{XVII} is a hydrogen atom; a linear or branched alkyl group having 1 to 8 carbon atoms, a cycloalkyl group having 3 to 10 carbon atoms, a cycloalkylalkyl group composed of a cycloalkyl moiety having 3 to 10 carbon atoms and an alkyl moiety having 1 to 3 carbon atoms, a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group carrying an alkyl moiety having 1 to 4 carbon atoms;

R⁵_{XVII} is a hydrogen atom, an alkyl group having 1 to 6 carbon atoms, a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group carrying an alkyl moiety having 1 to 4 carbon atoms; and

R⁶_{XVII} is an alkyl group having 1 to 6 carbon atoms, an alkenyl group having 3 to 6 carbon atoms, an alkynyl group having 3 to 6 carbon atoms or a substituted or unsubstituted aryl group.

[0228] Preferable examples of the compounds represented by the general formula (XVII) are those satisfying the fol-

lowing requirements:

5

10

20

25

- (1) A compound wherein mXVII is 5 and R1, R2 and R3 are each a hydrogen atom.
- (2) A compound wherein R⁴_{XVII} is a cycloalkyl group, such as monocycloalkyl, bicycloalkyl and tricycloalkyl groups. A preferable example of the monocycloalkyl group is a cyclohexyl group. A preferable example of the bicycloalkyl group is a norbornyl group, more preferably a 2-exo-norbornyl group. A preferable example of the tricycloalkyl group is an adamantyl group, more preferably a 1-adamantyl group.
 - (3) A compound wherein R⁴_{XVII} is a substituted or unsubstituted phenyl group or a substituted or unsubstituted phenylalkyl group.

 - (4) A compound wherein R⁵_{XVII} is a hydrogen atom.
 (5) A compound wherein A^{XVII} is S and R⁶_{XVII} is a lower alkyl group.
 - (6) A compound wherein a lower alkyl group is a methyl group.

[0229] R¹ and R² are preferably selected as specified for the formula (A).

[0230] According to a eighteenth aspect, the invention is directed to non imidazole compounds having the following formula (XVIII), analogous to those disclosed in Van der Goot et al. (Eur. J. Med. Chem. (1992) 27, 511-517):

$$\begin{array}{c} R^{t} \\ N \longrightarrow (CH_{2})_{t_{XVIII}} \longrightarrow S \longrightarrow C \longrightarrow NH \longrightarrow (CH_{2})_{UXVIII} \longrightarrow \\ R^{f}_{XVIII} \longrightarrow (XVIII) \end{array}$$

in which:

- R¹ and R² are as defined with reference to formula (A);
- R^eXVIII is H, alkyl or cycloalkyl;
- R^f_{XVIII} is H or halogen, in particular Cl, F, Br, or an alkyl;
 - t_{XVIII} is 1 to 3;
 - u_{XVIII} is 1 to 4.
 - [0231] Preferred groups R¹ and R² are as defined with reference to formula (A).
- [0232] Representative example is compound 83.
 - [0233] According to the invention, the W residue as defined in formula (A) and in particular as illustrated by formulae (I) to (XVIII), preferably contains no imidazole moiety attached in 4(5)-position and more preferably W contains no imidazole moiety.
 - [0234] The compounds according to the invention may be prepared according to one of the following schemes:
- [0235] More specifically, compounds of formula (I) can be obtained by the schemes 1 to 5:
 - [0236] In these schemes, R1, R2, R3, X and n are as defined in general formula (I).
 - [0237] Me and Et are intended to mean methyl and ethyl.

55

45

SCHEME 1 (methods A, B, C, D, H and K):

$$(R^{3})_{n_{3}} \xrightarrow{BrC_{n}H_{2n}Br} (R^{3})_{n_{3}} \xrightarrow{HNR^{1}R^{2}} (R^{3})_{n_{3}} \xrightarrow{XC_{n}H_{2n}NR^{1}R^{2}} (R^{3})_{n_{3}} \xrightarrow{XC_{n}H_{2n}NR^{1}R^{2}} (R^{3})_{n_{3}} \xrightarrow{R^{3}} (R^{3})_{n_{3}} \xrightarrow{R^{3}} (R^{3})_{n_{3}} \xrightarrow{R^{3}} (R^{3})_{n_{3}} \times (R^{3})_{n_{3}}$$

SCHEME 2 (methods F and L):

$$(R^{3})_{n_{3}} + HOC_{n}H_{2n}NR^{1}R^{2} \xrightarrow{R'OOC-N=N-COOR'} (R^{3})_{n_{3}} + OC_{n}H_{2n}NR^{1}R^{2} \xrightarrow{CC_{\theta}H_{5})_{3}P, THF, N_{2}} (R^{3})_{n_{3}} + OC_{n}H_{2n}NR^{1}R^{2}$$

SCHEME 3 (method E):

SCHEME 4 (method G):

$$H_3C$$

$$CH$$

$$CH$$

$$Et_2O$$

$$OC_nH_{2n}NR^1R^2$$

$$OC_nH_{2n}NR^1R^2$$

$$OC_nH_{2n}NR^1R^2$$

SCHEME 5 (method J):

$$H_3C$$

$$H_2NOH; HCI$$


$$MeOH and H_2O$$


$$OC_nH_{2n}NR^1R^2$$

$$OC_nH_{2n}NR^1R^2$$

$$OC_nH_{2n}NR^1R^2$$


(c) KI, K₂CO₃, EtOH, 3 h, 60 °C; (d) 6N HCI, 2 h, 100 °C.





For example: (a) dioxane/H₂O (1+1), 4 h, 0 °C; (b) acetonitrile, 5min, r.t.; (c) N-Boc-diphenylimido carbonate, 10 h, reflux; (d) 1N HCl, 0.5 h, reflux.


p, q, r = 0 - 3 (independently); t = 2 - 5.





 $R^{*d} = H \text{ or lower alkyl}$; $R^5 = alkyl$, cycloalkyl, or aryl.

p, q, r = 0 - 3 (independently); t = 2 - 5.

For example: (a) toluene, 12 h, r.t.; (b) toluene, tetrabutylammonium iodide, 15-crown-5, 12 h, 80 °C; (c) acetonitrile, 4 (d) thionyl chloride, THF, 12 h, 50 °C; (e) K₂CO₃, H₂O, EtOH, 2 days, reflux.








 $R^{e-d} = H$ or lower alkyl.

p, q, r = 0 - 3 (independently); t = 2 - 5.

For example: (a) acetone, triethylamine, 8 h, 50 °C; (b) NaH, MeOH, DMF, 6 h, 80 °C.

SCHEME 11:



 $R^{a-d} = H$ or lower alkyl; $R^5 = alkyl$, cycloalkyl, or aryl.

p, q, r = 0 - 3 (independently); t = 0 - 2 (independently).

For example: (a) toluene, 100 °C, nitrogen atmosphere, 12 h; (b) MeOH, SOCl₂; (c) triethylamine, MeOH.

SCHEME 12

$$\begin{pmatrix} R^{b}C \\ R^{c}C \\ R^{c}C \\ R^{c}C \\ R^{c}C \\ R^{c}C \\ C(CHR^{3})_{q} \end{pmatrix} = C(CH_{2})_{r} - N(CH_{2})_{r} -$$

 $R^{ad} = H$ or lower alkyl; X = Cl, Br, etc.

p, q, r = 0 - 3 (independently); t = 2 - 5; u = 1 - 5; w = 0 - 2.

(c) triethylamine, CH2Cl2, argon atmosphere, 0 °C, 18 h; (d) NaH, DMF, argon atmosphere, -15 °C; (e) 1N HCl, MeOH, 18 h, reflux. For example: (a) triethylamine, CH2Cl2, 24 h, r.t, (b) N,N,N',N'-tetramethylazodicarboxamide, tributylphosphine, MeOH, benzene, 24 h, r.t.;

ত্ Œ SORNETS

 $R^{a\cdot d} = H$ or lower alkyl; $X = NO_2$, NH_2 , OCH_3 , etc.

д q г=0-3(independently); t=2-6

For example: (a) triethylamine, CH₂Cl₂, 24 h, 50 °C; (b) triethylamine, KI, EtOH, 6 h, reflux; (c) thionyl chloride, THF, 2 h, 0 °C; (d) K₂CO₃, KI, EtOH, 6 h, reflux. SCHEME 14:

5

10

15

20

25

30

35

40

45

50

55

 $\begin{pmatrix}
R^{b}C \\
R^{c}C
\end{pmatrix}^{(CHR^{b})_{p}} N - CH_{(2)} - CH_{(2)} - C - OCH_{3}$ $(a) \qquad So_{2} - CH_{2} - CH_{2} - (CH_{2})_{u} - R$ $(b) \qquad NaH$ $(c) \qquad Sml_{2}$

$$\begin{pmatrix}
R^{b}C \\
R^{c}C
\end{pmatrix}^{(CHR^{a})_{p}} N - CH_{(2)} - CH_{(2)} - C = C - CH_{2} - (CH_{2})_{u} - R^{5}$$

$$(d) \text{ or } (d')$$

 $\begin{pmatrix}
R^{b}C \\
R^{c}C
\end{pmatrix}^{(CHR^{b})_{p}} N - CH_{(2)} - CH_{(2)} - HC = CH - CH_{2} - (CH_{2})_{u} - R^{5}$ (e)

$$\begin{pmatrix} R^{b}C \\ R^{c}C \\ \end{pmatrix}^{(CHR^{a})_{p}} N - CH_{(2)} - CH_{(2)} - CH_{2} - CH_{2} - CH_{2} - (CH_{2})_{u} - R^{6}$$

$$(CH_{2})_{t}$$

$$(CH_{2})_{t}$$

 $R^{a-d} = H$ or lower alkyl; $R^5 = alkyl$, cycloalkyl, or aryl. p, q, r = 0 - 3 (independently); t, u = 0 - 3.

For example: (a) n-BuLi, -78 °C; (b) THF, CIP(O)OEt₂; (c) THF, 4 mole% HMPA; (d) H₂, quinoline, ethyl acetate (cis); (d') Na/NH₃ (trans); (e) H₂, Pd (black), MeOH.

[0238] Detailed synthesis procedures are given in the examples.

[0239] The compounds of formula (A) according to the invention have antagonistic and/or agonistic properties at the

histamine H₃-receptors. They affect the synthesis and release of histamine monoamines or neuropeptides in brain and peripheral tissues.

- [0240] This property makes the compounds of the invention useful derivatives in human or veterinary medicine.
- [0241] Their therapeutical applications are those known for H₃-antagonist and/or agonist compounds and especially relate to the central nervous system disorders.
- **[0242]** Regarding antagonistic activity, the compounds according to the invention can be used in the treatment of Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.
- [0243] Regarding agonistic activity, the compounds according to the invention can be used in the treatment of various allergic and inflammatory diseases and as a sedative agent.
 - [0244] Therefore, the compounds of formula (A) according to the invention are advantageously used as active ingredient of medicaments which act as an antagonist and/or agonist of H_3 -receptors of histamine.
 - **[0245]** The antagonists are advantageously used as active ingredient in particular, of medicaments having psychotropic effects, promoting wakefullness, attention, memory and improving mood, in treatment of pathologies such as Alzheimer disease and other cognitive disorders in aged persons, depressive or simply asthenic states.
 - [0246] Their nootropic effects can be useful to stimulate attention and memorization capacity in healthy humans.
- [0247] In addition, these agents can be useful in treatment of obesity, vertigo and motion sickness.
- [0248] It can also be useful to associate the compounds of the invention with other psychiatric agents such as neuroleptics to increase their efficiency and reduce their side effects.
- [0249] Application in certain form of epilepsy is also foreseen.
 - [0250] Their therapeutic applications involve also peripheral organs mainly a stimulant of secretions or gastro-intestinal motricity.
 - [0251] The compounds of the invention are particularly useful for the treatment of CNS disorders of aged persons.
 - [0252] The agonists are advantageously used as active ingredients in the treatment of acute and chronic inflammatory diseases such as arthritis, cystitis, bronchitis, asthma and other pneumopathies, inflammatory bowel diseases such as Crohn disease.
 - [0253] Their inhibitory effect on sensory fibers can be useful to depress excessive vegetative reflexes such as in urinary incontinence, to prevent nociceptive sensations and treat pruritus.
 - [0254] Their inhibitory effect on sympathetic and peptidergic fibers can be useful in cardiovascular diseases such as coronary or vascular diseases in which their activity is excessive. Their inhitory effect on the activity of enterochromaffin or enterochromaffin-like cells can be useful in gastro-intestinal hypersecretion manifestations.
 - [0255] Their inhibitory effect on histaminergic and other aminergic neurons in the central nervous system can be useful in the treatment of sleep disorders or as sedative agents in various psychiatric manifestations.
 - **[0256]** The present invention also relates to medicaments having the above-mentioned effects comprising as active ingredient, a therapeutically effective amount of a compound of formula (A).
 - [0257] The present invention relates more particularly to such medicaments containing a compound of formula (I) to (XVIII).
 - [0258] The present invention also relates to pharmaceutical compositions containing as active ingredient a therapeutically effective amount of a compound (A) together with a pharmaceutically acceptable vehicle or excipient.
- [0259] The invention is directed to such pharmaceutical compositions containing as active-ingredient, a compound of formula (I) to (XVIII).
 - [0260] The medicaments or pharmaceutical compositions according to the invention can be administered via oral, parenteral or topical routes, the active ingredient being combined with a therapeutically suitable excipient or vehicle.
 - [0261] According to the invention, oral administration is advantageously used.
 - [0262] Another subject of the present invention is the use of the compounds of formula (A) for the preparation of H₃-antagonist and/or agonist medicaments according to the above-mentioned forms.
 - [0263] The invention further relates to the use of the compounds of formula (A) for preparing medicaments having the pre-cited effects.
 - [0264] The invention also concerns the use of a compound of formula (I) to (XVIII).
- [0265] Still another subject of the invention is a method for the treatment of precited ailments comprising administering a therapeutically effective dose of a compound (I), optionally in combination with a therapeutically acceptable vehicle or excipient.
 - [0266] The invention is also directed to such a method comprising administering a therapeutically effective dose of a compound of formula (I) to (XVIII).
- 5 [0267] For each of the above-indications, the amount of the active ingredient will depend upon the condition of the patient.
 - [0268] However, a suitable effective dose will be in general in the range of from 10 to 500 mg per day and of from 1 to 10 mg/day for particularly active compounds.

[0269] These doses are given on the basis of the compound and should be adapted for the salts, hydrates or hydrated salts thereof.

[0270] The invention is now illustrated by the following examples.

5 EXAMPLES

[0271] The structure of the synthesized compounds and their method of preparation as well as their melting point, recrystalisation solvant and elemental analysis are summarized in the following Table I:

TABLE 1:

4	5		

И	FORMULA	mp	analysis (calc.)	method
- /	STRUCTURE	(recryst. solv)		
	NAME			
1	C ₁₆ H ₂₅ NO; C ₂ H ₂ O ₄	143-145°C	C: 64.06 (64.07)	A
		(absolute ethanol)	H: 8.09 (8.16)	Ì
	O-(CH 2)s-N (COOH) 2		N: 4.14 (4.15)	
	1-(5-phenoxypentyl)-piperidine hydrogen oxalate			<u> </u>
2	C ₁₅ H ₂₃ NO; C ₂ H ₂ O ₄	153-155°C	C: 63.06 (63.14)	A
		(absolute ethanol)	H: 7.78 (7.79)	
	O-(CH 2)5-N (COOH) 2		N: 4.42 (4.33)	
	1-(5-phenoxypentyl)-pyrrolidine hydrogen oxalate	}	·	<u> </u>
3	C14H23NO; C2H2O4	122-124°C	C: 61.74 (61.72)	A
	CH ₃ (COOH)	(absolute ethanol)	H: 8.24 (8.09)	}
l	O-(CH ₂) ₅ -N CH ₂ CH ₃ (COOH) ₂		N: 4.52 (4.50)	
	N-methyl-N-(5-phenoxypentyl)-ethylamine hydrogen]
	oxalate			<u> </u>
4	C ₁₅ H ₂₃ NO ₂ ; C ₂ H ₂ O ₄	166-168°C	C: 60.10 (60.16)	A
		(absolute ethanol)	H: 7.45 (7.31)	
	O-(CH 2)5N O (COOH) 2		N: 4.08 (4.13)	
	1-(5-phenoxypentyl)-morpholine hydrogen oxalate			
5	C ₁₇ H ₂₇ NO; C ₂ H ₂ O ₄	132-134°C	C: 64.70 (64.93)	A
ł		(absolute ethanol)	H: 8.34 (8.32)	į
	O-(CH 2)5 N (COOH) 2		N: 3.85 (3.99)	
	N-(5-phenoxypentyl)-hexamethyleneimine hydrogen	\		
	oxalate			
6	C ₁₆ H ₂₇ NO; C ₂ H ₂ O ₄	90-91°C	C: 63.60 (63.69)	В
		(isopropyl alcohol)	H: 8.81 (8.61)	
	O-(CH 2)5-N (COOH) 2		N: 3.97 (4.13)	
1	N-ethyl-N-(5-phenoxypentyl)-propylamine hydrogen		1	
L	oxalate	<u> </u>	<u> </u>	

r					
ŧ	7	C ₁₇ H ₂₇ NO; 1.1 C ₂ H ₂ O ₄	80-83°C	C: 64.15 (63.98)	В
5		į	(isopropyl alcohol)	H: 8.42 (8.17)	
	ł	CH ₃	ĺ	N: 3.97 (3.89)	
		-0-(CH ₂) ₅ -N 1.1 (СООН) 2		į	- 1
	- 1	G (3/12/3)		İ	1
10	İ	1-(5-phenoxypentyl)-2-methyl-piperidine hydrogen			1
		oxalate			
	8	C19H31NO; C2H2O4	165-166°C	C: 66.27 (66.46)	В
	1		(absolute ethanol)	H: 8.94 (8.76)]
15	ľ		(N: 3.72 (3.69)	1
		O-(CH 2)5-N nC 3 H7 (COOH) 2		11. 3.72 (3.07)	
		1-(5-phenoxypentyl)-4-propyl-piperidine hydrogen			- 1
20		oxalate			l
20	9	C ₁₇ H ₂₇ NO; C ₂ H ₂ O ₄	151-152°C	C: 64.87 (64.93)	В
		O-(CH ₂) ₅ -N CH ₃ (COOH) ₂	(absolute ethanol)	H: 8.41 (8.32)	1
				N: 4.01 (3.99)	I
25		1-(5-phenoxypentyl)-4-methyl-piperidine hydrogen			
		oxalate			
	10	C ₁₇ H ₂₇ NO; C ₂ H ₂ O ₄	140-141°C	C: 65.35 (64.93)	В
,]	_у сн _з	(isopropyl alcohol)	H: 8.49 (8.32)	
30		O-(CH ₂) ₆ N (COOH) ₂		N: 4.00 (3.99)	1
		0 -(ch 2/6 H) (cost) 2			1
		1-(5-phenoxypentyl)-3-methyl-piperidine hydrogen	}	•	1
I		oxalate			}
35	11	C ₁₇ H ₂₆ N ₂ O ₂ ; C ₂ H ₂ O ₄	186-188°C	C: 59.78 (59.99)	В
	'''		(absolute ethanol)	Н: 7.47 (7.42)	_
		O-(CH ₂) ₅ -N NCOCH ₃ (COOH) ₂	(2000,200	N: 7.35 (7.36)	
	}	1-acetyl-4-(5-phenoxypentyl)-piperazine hydrogen		2 7 7 7 7	
40	}	oxalate			
	12	C ₁₈ H ₂₉ NO; 1.05 C ₂ H ₂ O ₄	154-155°C	C: 65.16 (65.25)	В
	"	7102-27	(absolute ethanol)	H: 8.61 (8.47)	-
45	{	ECH 3	(absolute ethalibi)	N: 3.66 (3.79)	}
40	1			N: 3.00 (3.79)	{
		O -(CH 2)5-N 1.05 (COOH) 2	1	1	
	1				
50	1	CH ₃	1		
	1	1-(5-phenoxypentyl)-3,5-trans-dimethyl-piperidine	1		
	L	hydrogen oxalate		<u> </u>	

ſ					
	13	C ₁₈ H ₂ 9NO; C ₂ H ₂ O ₄	154-155°C	C: 65.62 (65.73)	В
5			(isopropyl alcohol)	H: 8.64 (8.55)	
		CH ₃		N: 3.63 (3.83)	1
		. (COOH)2			}
		5 (0.123.)			ŀ
10	i	CH₃			
		1-(5-phenoxypentyl)-3,5-cis-dimethyl-piperidine			
		hydrogen oxalate			
	14	C ₁₈ H ₂₉ NO; HCl	135-136°C	C: 69.18 (69.32)	В
15			(acetone)	H: 9.79 (9.70)	į
		сн,		N: 4.28 (4.49)	[
		O-(CH 2)5'N HCI			
)
20		CH 3			
		1-(5-phenoxypentyl)-2,6-cis-dimethyl-piperidine			
	<u> </u>	hydrochloride			
	15	C ₁₉ H ₂₉ NO ₃ ; C ₂ H ₂ O ₄	149-150°C	C: 61.16 (61.60)	В
25			(absolute ethanol)	H: 7.76 (7.63)	į
	1		ļ	N: 3.40 (3.42)	1
	l	Сос 2H ₅ (соон) 3	1		
30	1	4-carboethoxy-1-(5-phenoxypentyl)-piperidine			
00	1	hydrogen oxalate			
	16	C ₁₉ H ₂₉ NO ₃ ; C ₂ H ₂ O ₄	117-118°C	C: 61.54 (61.60)	В
] "		(isopropyl alcohol)	Н: 7.87 (7.63)]
35	1	,cooc ₂H ₆	(isopropy: alcohol,	N: 3.29 (3.42)	
	1				
		O -(CH 2)5'N (COOH) 2			
	•	3-carboethoxy-1-(5-phenoxypentyl)-piperidine			
40		hydrogen oxalate			
	17	C ₁₆ H ₂₃ NO; C ₂ H ₂ O ₄	177-179°C	C: 64.19 (64.46)	В
	\ '	5101-251-15, 521-254	(methanol)	H: 7.49 (7.51)	-
	1		(titemanor)	N: 4.25 (4.18)	
45	1	O-(CH ₂) ₆ N (COOH) ₂		14: 4.23 (4.10)	
	1				
		1-(5-phenoxypentyl)-1,2,3,6-tetrahydropyridine			
		hydrogen oxalate	1	L	<u> </u>
50					

Ī		C ₁ 5H ₂₂ N ₂ O ₃ ; C ₂ H ₂ O ₄ ; 0.2 H ₂ O	145-147°C	C: 54 90 (54 90)	С
	18	015112211203, 0211204, 012 1120	(absolute ethanol)	C: 54.89 (54.89)	٠ <u> </u>
5	1	(COOH) 2	(absolute emanoi)	H: 6.68 (6.61)	ì
		02N-()-0-(CH 2)5N.		N: 7.41 (7.53)	1
		0.2 H ₂ O	i,	ļ	1
-		1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine hydrogen			ł
10		oxalate			
	19	C ₁₅ H ₂₂ CINO; C ₂ H ₂ O ₄	139-141°C	C: 57.00 (57.06)	С
			(absolute ethanol)	H: 6.63 (6.76)	1
		CI-(CH ₂) ₅ -N (COOH) ₂		N: 3.79 (3.91)	
15				C1: 10.24 (9.91))
		1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine hydrogen			1
	ļ	oxalate			
20	20	C ₁₆ H ₂₅ NO ₂ ; C ₂ H ₂ O ₄	115-116°C	C: 61.22 (61.17)	С
		_	(absolute ethanol)	H: 7.72 (7.70)	
!		H ₃ CO-(CH ₂) ₅ N (COOH) ₂		N: 4.03 (3.96)	
		11,500		į	1
25		1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine			
		hydrogen oxalate			
	21	C ₁₆ H ₂₅ NO; C ₂ H ₂ O ₄	138-140°C	C: 64.05 (64.07)	С
	İ		(absolute ethanol)	H: 8.00 (8.07)	
30		H ₃ C-(CH ₂) ₅ N (COOH) ₂		N: 4.10 (4.15)	
		1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine hydrogen	}		}
35	}	oxalate			
JJ	22	C ₁₆ H ₂₂ N ₂ O; 1.1 C ₂ H ₂ O ₄	129-130°C	C: 61.24 (61.16)	С
			(absolute ethanol)	Н: 6.81 (6.82)	
	}	A		N: 7.95 (7.84)	
40	1	NC-(CH 2)5N 1.1 (COOH) 2			
	}	1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine hydrogen			
	1	oxalate	\		(
	23	C ₁₉ H ₂ 5NO; C ₂ H ₂ O ₄	166-167°C	C: 67.42 (67.54)	С
45	1		(methanol)	Н: 7.26 (7.29)	
	1		,,	N: 3.66 (3.75)	
	1	O-(CH 2) 5-N (COOH) 2			1
50	1				
3 <i>0</i>	i	1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine hydrogen			
	1	oxalate			
					

				
24	C ₁₉ H ₂₅ NO; 1.25 C ₂ H ₂ O ₄	160-163°C	C: 65.12 (65.22)	С
1 1		(methanol)	H: 7.17 (7.00)	. }
	O-(CH 2)5N 1.25 (COOH) 2		N: 3.52 (3.54)	
	1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine hydrogen	į.		1
	oxalate	1		j
25	C ₁₅ H ₂₂ CINO; C ₂ H ₂ O ₄	131-132°C	C: 56.94 (57.06)	С
		(absolute ethanol)	H: 6.67 (6.76)	1
	CI		N: 3.74 (3.91)	1
	O-(CH 2)5-N (COOH) 2		Cl: 9.64 (9.91)	
	1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidiae hydrogen oxalate			
26	C ₂₁ H ₂ 7NO; C ₂ H ₂ O ₄	189-190°C	C: 69.16 (69.15)	С
20		(methanol)	H: 7.39 (7.32)	Ĭ
	O-(CH 2)5-N (COOH) 2	(medianol)	N: 3.39 (3.51)	
	1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine hydrogen			. 1
	oxalate			
27	C ₁₉ H ₂₉ NO; C ₂ H ₂ O ₄	131-132°C	C: 66.73 (66.82)	С
1		(absolute ethanol)	H: 8.37 (8.28)	
	O-(CH 2) 5N (COOH) 2		N: 3.68 (3.71)	
1			•	
	1-{5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-		İ	
<u></u>	pyrrolidine hydrogen oxalate			ļ
28	C ₂₁ H ₂₇ NO; 1.1 C ₂ H ₂ O ₄	155-157°C	C: 68.40 (68.22)	С
		(absolute ethanol)	H: 7.04 (7.21)	1
	-0-(CH 2)5N 1.1 (COOH) 2		N: 3.45 (3.43)	}
	1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine hydrogen			
	oxalate		<u> </u>	

ſ	29	C ₁₅ H ₂₁ NO; C ₂ H ₂ O ₄	140-141°C	C: 63.45 (63.54)	В
_ }	}		(absolute ethanol)	H: 7.26 (7.21)	j
5		O-(CH 2)5N (COOH) 2		N: 4.26 (4.36)	
	Ì	1-(5-phenoxypentyl)-2,5-dihydropyrrole hydrogen			1
10		oxalate			
	30	C ₁₉ H ₂₉ NO; C ₂ H ₂ O ₄	148-149°C	C: 66.99 (66.82)	С
	1	ì	(absolute ethanol)	H: 8.47 (8.28)	1
15		O-(CH ₂) ₅ -N (COOH) ₂		N: 3.72 (3.71)	
	1	1-{5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-	1		!
20		руттоlidine hydrogen oxalate			
	31	C14H21NO; C2H2O4	143-144°C	C: 62.25 (62.12)	С
			(absolute ethanol)	H: 7.46 (7.49)	
25		O-(CH 2) (COOH) 2		N: 4.49 (4.53)	
		1-(4-phenoxybutyl)-pyrrolidine hydrogen oxalate			
	32	C ₁₆ H ₂₅ NO; 1.1 C ₂ H ₂ O ₄	146-147°C	C: 63.06 (63.10)	С
30			(absolute ethanol)	H: 8.03 (7.91)	
	 	O-(CH 2) g N 1.1 (COOH) 2		N: 4.32 (4.04)	
		1-(6-phenoxyhexyl)-pyrrolidine hydrogen oxalate			
35	33	C ₁₅ H ₂₃ NS; 1.1 C ₂ H ₂ O ₄	150-152°C	C: 59.52 (59.29)	С
)		(absolute ethanol)	H: 7.44 (7.29)]]
40		8-(CH ₂) ₅ -N 1.1 (COOH) ₂		N: 4.06 (4.02)	
		1-(5-phenylthiopentyl)-pyrrolidine hydrogen oxalate	<u>c</u>		
	34	C ₁₄ H ₂₁ NS; C ₂ H ₂ O ₄	114-116°C	C: 59.24 (59.05)	С
		1	(absolute ethanol)	H: 7.16 (7.12)) 1
45	1	(COOH) 2		N: 4.16 (4.30)	
		S-(CH 2)4N (COOH) 2		S: 9.79 (9.85)	
		1-(4-phenylthiobutyl)-pyrrolidine hydrogen oxalate			

T		T T		
35	C ₁₃ H ₁₉ NO; C ₂ H ₂ O ₄	169-170℃	C: 60.98 (61.00)	C
		(absolute ethanol)	H: 7.14 (7.17)	
	O-(CH 2)3TN (COOH) 2		N: 4.64 (4.74)	
	1-(3-phenoxypropyl)-pyrrolidine hydrogen oxalate			
36	C ₁₅ H ₂₂ N ₂ O ₃ ; C ₂ H ₂ O ₄	130-131°C	C: 55.30 (55.43)	C
1	02N	(absolute ethanol)	H: 6.55 (6.57)	
1	O-(CH ₂) ₅ -N (COOH) ₂		N: 7.49 (7.60)	
1	1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine hydrogen			
]	oxalate			
37	C ₁₅ H ₂₂ FNO; C ₂ H ₂ O ₄	149-150°C	C: 59.52 (59.81)	С
		(absolute ethanol)	H: 7.12 (7.09)	
	(COOH) 2		N: 4.05 (4.10)	
1 1	F-(CH ₂) ₅ -N (COOH) ₂			
1	1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine hydrogen			1
]]	oxalate			
38	C ₁₇ H ₂₆ N ₂ O ₃ ; C ₂ H ₂ O ₄	148-149°C	C: 57.32 (57.55)	С
		(absolute ethanol)	H: 7.19 (7.12)	
]	CH ₃		N: 6.89 (7.07)	}
	O ₂ N — (COOH) ₂		·	
	1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine			
	hydrogen oxalate			
39	C ₁₇ H ₂₅ NO ₂ ; C ₂ H ₂ O ₄	130-134°C	C: 62.43 (62.45)	D
Ì		(absolute ethanol)	H: 7.41 (7.45)	ĺ
	CH ₃ -C-(CH ₂) ₅ -N (COOH) 2		N: 3.75 (3.83)	
	1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine hydrogen			
-	oxalate		- 42 42 42 53	│
40	C ₁₅ H ₂₄ N ₂ O; 2.1 C ₂ H ₂ O ₄	120-122°C	C: 52.49 (52.72)	E1
1		(absolute ethanol)	1	
	H ₂ N-(CH ₂) ₅ N 2.1 (COOH) ₂		N: 6.32 (6.40)	
	1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine	1		
1	di-(hydrogen oxalate)	1	<u> </u>	1

ſ		C. Mark O. Callan		T	
	41	C ₁₆ H ₂₂ N ₂ O; C ₂ H ₂ O ₄	119-120°C	C: 61.95 (62.05)	C
5		NC	(absolute ethanol)	H: 6.88 (6.94)	Ĭ
]	1			N: 8.00 (8.04))
		O-(CH 2)5N (COOH) 2			1
10		1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine hydrogen]	}
[oxalate			
	42	C ₁₃ H ₂₀ N ₂ O ₃ ; C ₂ H ₂ O ₄	160-161°C	C: 52.46 (52.63)	F
			(absolute ethanol/	H: 6.49 (6.48)	j
15		CH ₂ CH ₃	methanol	N: 8.10 (8.12)	1
		O ₂ N — (COOH) ₂ (COOH) ₂	1:1)		
į			,	,	I
		N-[3-(4-nitrophenoxy)-propyl]-diethylamine			}
20		hydrogen oxalate			
	43	C ₁₄ H ₂₀ N ₂ O; C ₂ H ₂ O ₄	148-150°C	C: 59.40 (59.62)	F
			(absolute ethanol)	H: 6.82 (6.88)	
	1	NC-(CH 2)3N (COOH) 2		N: 8.60 (8.69)	
25		CH 2 CH 3			
		N-[3-(4-cyanophenoxy)-propyl]-diethylamine			ŀ
		hydrogen oxalate			l
	44	C ₂₂ H ₂ 7NO ₂ ; C ₂ H ₂ O ₄	141-142°C	C: 67.17 (67.42)	
30	***	022.121.102, 02.1204		C: 67.17 (67.43)	D
			(absolute ethanol)	H: 6.80 (6.84)	1
	İ	O -(CH 2)5-N (COOH) 2	,	N: 3.18 (3.28)	1
					Ì
35	Ì	1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine			
		hydrogen oxalate			
	45	C23H29NO2; C2H2O4	177-178°C	C: 67.77 (68.01)	D
40			(absolute ethanol)	H: 7.09 (7.08)	
40			}	N: 3.26 (3.17)]]
	1	CH ₂ ·C-CH ₂ ·C-CH ₂) ₅ -N	}	1	
		(COOH) 2	Į.		
45	l	1-{5-[4-(phenylacetyl)-phenoxy]-pentyl}-pyrrolidine		}	
	1)
	<u> </u>	hydrogen oxalate			لــــــل

				
46	C ₁₅ H ₂₃ NO ₂ ; 1.1 C ₂ H ₂ O ₄	108-110°C	C: 59.30 (59.30)	F
		(absolute ethanol)	H: 7.47 (7.29)	
	H ₃ C-C-C-C-C-C-C-C-C-C-C-C-C-C-C-C-C-C-C-		N: 4.18 (4.02)	
	N-[3-(4-acetylphenoxy)-propyl]-diethylamine		}	
	hydrogen oxalate			
47	C ₁₇ H ₂₆ N ₂ O ₂ ; C ₂ H ₂ O ₄	142-144°C	C: 59.67 (59.99)	С
		(absolute ethanol)	H: 7.55 (7.42)	
	H ₃ C-C-N O-(CH ₂) ₅ -N (COOH) ₂		N: 7.25 (7.36)	
	1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine			
	hydrogen oxalate			
48	C ₂₁ H ₂₇ NO ₂ ; C ₂ H ₂ O ₄	135-136°C	C: 66.49 (66.49)	D
		(absolute ethanol)	H: 7.05 (7.04)	
	O-(CH 2)5N (COOH) 2		N: 3.24 (3.37)	
}	1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine			
<u> </u>	hydrogen oxalate			
49	C ₂₂ H ₂₈ N ₂ O ₂ ; 1.1 C ₂ H ₂ O ₄	176-178°C	C: 64.56 (64.38)	E2
1		(absolute ethanol)	H: 6.89 (6.74)	
	C-N-C-N-CH 2)5N		N: 6.26 (6.20)	
•	1.1 (COOH) ₂			
١.	1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine			
	hydrogen oxalate			
50	C ₁₇ H ₂₇ NO ₂ ; C ₂ H ₂ O ₄	102-104°C	C: 61.89 (62.11)	G
l		(absolute ethanol)	Н: 7.94 (7.96)	l
	H ₃ C, HO CH-(CH ₂) ₅ N (COOH) ₂		N: 3.77 (3.81)	
	1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-			
	pyrrolidine hydrogen oxalate			
51		120-122°C	C: 61.56 (61.70)	н
		(absolute ethanol)	H: 7.54 (7.48)	1
	NC-(CH ₂) ₅ -N CH ₂ CH ₃ (COOH) ₂		N: 7.87 (7.99)	
	N-[5-(4-cyanophenoxy)-pentyl]-diethylamine			
1	hydrogen oxalate		<u> </u>	

1					
	52	C ₁₇ H ₂₄ N ₂ O; C ₂ H ₂ O ₄	115-116℃	C: 62.62 (62.97)	н
5			(absolute ethanol)	H: 7.20 (7.23)	1
Ĭ		NC-(CH 2)5N (COOH) 2		N: 7.76 (7.73)	1
		NC (CA 2)5 M (SSSA) 2		,	1
		1-[5-(4-cyanophenoxy)-pentyl]-piperidine hydrogen			1
10		oxalate	_	•	1
	53	C14H20N2O; C2H2O4	148-149°C	C: 59.68 (59.62)	н
			(absolute ethanol)	H: 6.76 (6.88)	
		CH3 (2001)	`	N: 8.57 (8.69)	. [
15		NC-(CH 2)5-N (COOH) 2			[
		NES (A company control of the state of the			. 1
		N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine			1
		hydrogen oxalate			
20	54	C ₁₃ H ₁₈ N ₂ O; C ₂ H ₂ O ₄	124-125°C	C: 58.15 (58.43)	H
		CH 2CH 3	(absolute ethanol)	H: 6.30 (6.54)	
		NC-(CH 2) 2N (COOH) 2 CH 2CH 3		N: 8.95 (9.09)	
		CH ₂ CH ₃			1
25		N-[2-(4-cyanophenoxy)-ethyl]-diethylamine hydrogen			1
:		oxalate			
	55	C ₁₂ H ₁₆ N ₂ O; C ₂ H ₂ O ₄	166-167°C	C: 57.01 (57.14)	н
30		<u></u>	(absolute ethanol/	H: 6.02 (6.16)	l
		NC-(CH 2)3-N (COOH) 2	methanol	N: 9.46 (9.52)	
		CH ²	1:1)]
		N-[3-(4-cyanophenoxy)-propyl]-dimethylamine			
35		hydrogen oxalate			I
	56	C ₁₅ H ₂₂ N ₂ O; C ₂ H ₂ O ₄	143-145°C	C: 60.80 (60.70)	н
			(absolute ethanol)	H: 7.11 (7.19)	
	ł	CH ₂ CH ₃	[N: 8.22 (8.33)	
40	-	NC-(CH ₂) ₄ N (COOH) ₂ CH ₂ CH ₃		, ,	
	ĺ				
	l	N-[4-(4-cyanophenoxy)-butyl]-diethylamine hydrogen			
	<u> </u>	oxalate	 		
45	57	C ₁₈ H ₂₈ N ₂ O; C ₂ H ₂ O ₄	134-136°C	C: 63.38 (63.47)	н
					1 1
		C-H-	(absolute ethanol)	H: 8.11 (7.99)	}
		NC-(CH 2)5-N (COOH) 2	(absolute ethanol)	H: 8.11 (7.99) N: 7.29 (7.40)	
50		NC-(CH ₂) ₅ -N(C ₃ H ₇ (COOH) ₂	(absolute ethanol)	· · ·	
50			(absolute ethanol)	· · ·	
50		С ₃ Н _{7.}	(absolute ethanol)	· · ·	

	0 11 11 011 0		T	
58	C ₁₄ H ₁₈ N ₂ O; 1.1 C ₂ H ₂ O ₄	163-165°C	C: 58.95 (59.08)	н
{		(absolute ethanol)	H: 6.23 (6.18)	1
	NC-(CH 2)3-N 1.1 (COOH) 2		N: 8.43 (8.51)	
} {	1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine hydrogen		}	
	oxalate			
59	C ₁₅ H ₂₀ N ₂ O; 1.05 C ₂ H ₂ O ₄	151-153°C	C: 60.62 (60.61)	Н
1 1		(absolute ethanol)	H: 6.66 (6.57)	
	NC-(CH 2)31N 1.05 (COOH) 2		N: 8.25 (8.27)	
	1-[3-(4-cyanophenoxy)-propyl]-piperidine hydrogen oxalate			
60	C ₁₆ H ₂₂ N ₂ O; 1.05 C ₂ H ₂ O ₄	124-125°C	C: 61.62 (61.60)	н
		(absolute ethanol)	H: 6.94 (6.88)	
			N: 7.87 (7.94)	
	NC-(CH 2)3-N 1.05 (COOH) 2			
	N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine	1		
<u></u>	hydrogen oxalate			
61	C ₁₇ H ₂₆ N ₂ O; C ₂ H ₂ O ₄	110-112°C	C: 62.90 (62.62)	н
1		(absolute ethanol)	H: 7.76 (7.74)	
	NC-(CH ₂) ₆ -N, (COOH) ₂ .	·	N: 7.61 (7.69)	
}	N-[6-(4-cyanophenoxy)-hexyl]-diethylamine	}		
	hydrogen oxalate			
62	C ₁₆ H ₂₄ N ₂ O; C ₂ H ₂ O ₄	127-128°C	C: 61.57 (61.70)	н
1		(absolute ethanol)	H: 7.57 (7.48)	l
	NC-(CH 2)5-N (COOH) 2		N: 7.91 (7.99)	
}	N-[3-(4-cyanophenoxy)-propyl]-dipropylamine	}	}	ł
	hydrogen oxalate			
63	C ₁₅ H ₂₅ NO ₂ ; C ₂ H ₂ O ₄ ; 0.5 H ₂ O	33-36°C	C: 58.15 (58.27)	G
		(isopropyl alcohol)		1
1	H ₃ C. C ₂ H ₅ (COOH) ₂		N: 4.21 (4.00)	1
	HO CH- C2H5 0.5 H2O			
	N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-			
- 1	diethylamine hydrogen oxalate hemihydrate	Į.	1	1

5

64 C15H24N2O2; C2H2O4 99-100°C C: 57.26 (57.61) H ₃ C HO-N 4'-(3-diethylaminopropoxy)-acetophenone-oxime hydrogen oxalate C16H23NO2; C2H2O4 65 C16H23NO2; C2H2O4 159-160°C (absolute ethanol) 170 C: 57.26 (57.61) H: 7.47 (7.39) N: 7.72 (7.90) 180 C: 57.26 (57.61) H: 7.47 (7.39) N: 7.72 (7.90) 180 C: 61.18 (61.52) (absolute ethanol) H: 7.11 (7.17)	K.
10 H3C C2H5 C2H5 (COOH) 2 C2H5 (COOH) 2 C2H5 (COOH) 2 C2H5 (COOH) 2 C2H5 (COOH) 2 C2H5 (COOH) 2 C2H5 (COOH) 2 C16H23NO2; C2H2O4 (COOH) 2 C2H5 (COOH) 2 C2	K .
10 C ₂ H ₅ 4'-(3-diethylaminopropoxy)-acetophenone-oxime hydrogen oxalate C ₁₆ H ₂₃ NO ₂ ; C ₂ H ₂ O ₄ 159-160°C (absolute ethanol) H: 7.11 (7.17)	K .
10 hydrogen oxalate 65 C16H23NO2; C2H2O4 159-160°C C: 61.18 (61.52) (absolute ethanol) H: 7.11 (7.17)	K
65 C ₁₆ H ₂₃ NO ₂ ; C ₂ H ₂ O ₄ 159-160°C C: 61.18 (61.52) (absolute ethanol) H: 7.11 (7.17)	К.
(absolute ethanol) H: 7.11 (7.17)	K
	.
	1
15 H ₃ C-C-O-(CH ₂) ₃ -N (COOH) ₂ N: 3.96 (3.99)	- 1
1-[3-(4-acetylphenoxy)-propyl]-piperidine hydrogen	
oxalate	
20 66 C ₁₇ H ₂ 5NO ₂ ; C ₂ H ₂ O ₄ 143-144°C C: 62.11 (62.45)	ĸ
(absolute ethanol) H: 7.41 (7.45)	İ
CH ₃ N: 3.79 (3.83)	
H ₃ C-C (CH ₂) ₃ -N (COOH) ₂	ļ
25	
1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine	1
hydrogen oxalate	
30 67 C18H27NO2; C2H2O4 171-172°C C: 63.06 (63.31)	K
(absolute ethanol) H: 7.44 (7.70)	1
N: 3.64 (3.69)	1
H ₃ C-C-(CH ₂) ₃ -N (COOH) ₂	l l
35 H ₃ C-G-(CH ₂) ₃ -N (CDOH) ₂	- 1
CH ₃	i
1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-	· ·
piperidine hydrogen oxalate	
40 68 C ₁₇ H ₂₅ NO ₂ ; C ₂ H ₂ O ₄ 160-161°C C: 62.47 (62.45)	ĸ
(absolute ethanol) H: 7.46 (7.45)	1
H ₃ C-C-C-CH ₂) ₃ N CH ₃ (COOH) 2	
1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine	ľ
hydrogen oxalate	


69	C ₁₇ H ₂₅ NO ₂ ; C ₂ H ₂ O ₄	148-149°C	C: 62.54 (62.45)	L
		(absolute ethanol)	H: 7.51 (7.45)	l
	C2H5-C-(CH2)3-N (COOH) 2		N: 3.79 (3.83)	
	ے ۔			·
	1-[3-(4-propionylphenoxy)-propyl]-piperidine			Ì
<u> </u>	hydrogen oxalate			
70	C ₁₈ H ₂₇ NO ₂ ; C ₂ H ₂ O ₄	174-175°C	C: 63.22 (63.31)	ĸ
į		(absolute ethanol)	H: 7.60 (7.70)	
	CH3		N: 3.64 (3.69)	
	H ₃ C-G-(CH ₂) ₃ -N (COOH) ₂			
	CH ₃			
1	1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-			
 	piperidine hydrogen oxalate			
71	C ₁₅ H ₂₁ NO ₂ ; C ₂ H ₂ O ₄	152-153℃	C: 60,23 (60,52)	L
		(absolute ethanol)	H: 6.81 (6.87)	
	H-G-(CH ₂) ₃ -N (COOH) ₂		N: 4.15 (4.15)	
1	1-[3-(4-formylphenoxy)-propyl]-piperidine hydrogen			
L	oxalate			
72	C ₁₈ H ₂₇ NO ₂ ; C ₂ H ₂ O ₄	121-122°C	C: 63.02 (63.31)	L
		(absolute ethanol)	H: 7.73 (7.70)	l
	H ₃ C, CH -C-C-C-C-CH ₂) ₃ -N		N: 3.66 (3.69)	
Ì	(COOH) 2	l		
	1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine			1
	hydrogen oxalate		<u> </u>	<u> </u>
73	C ₁₆ H ₂₅ NO ₂ ; 1.5 C ₂ H ₂ O ₄	118-120°C	C: 57.27 (57.28)	L
		(absolute ethanol)	H: 7.00 (7.08)	
	C ₂ H ₅ -C-C-CH ₂) ₃ -N C ₂ H ₅ 1.5 (COOH) 2		N: 3.47 (3.52)	
1	Ö € C₂H₅	}		}
1	N-[3-(4-propionylphenoxy)-propyl]-diethylamine		1	
<u></u>	hydrogen oxalate	<u> </u>		<u> </u>

	74	C ₁₈ H ₂₇ NO ₂ ; C ₂ H ₂ O ₄	138-139°C	C: 63.09 (63.31)	L
5		C ₃ H ₇ -C-(CH ₂) ₃ N (COOH) ₂	(absolute ethanol)	H: 7.78 (7.70) N: 3.75 (3.69)	·
10		1-[3-(4-butyrylphenoxy)-propyl]-piperidine hydrogen oxalate			
	75	C ₁₆ H ₂₁ NO ₂ ; 1.1 C ₂ H ₂ O ₄	143-144°C (absolute ethanol)	C: 61.21 (61.00) H: 6.25 (6.52)	K
15		H3C-6-(CH 5)3-M		N: 4.00 (3.91)	
		1.1 (COOH) 2 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-			
20		tetrahydropyridine hydrogen oxalate			


_	40 E	<i>35</i>	30	25 Synthesis	20	15	10	5
NO	Sunciare							
	$\langle \rangle$	\Rightarrow		5	(в)	2 3		
7.1	$\langle \rangle$		\Diamond		HaN Ho'		\wp	
78	\(\)		<u></u>	\bigsig \bigsi	Hah Hah	H,C-80,	3	
79				\\	он Ман	2 3 3 S	\bigcirc	
80			$\boldsymbol{\wedge}$	8	O=C=IA			
26		_ ⊶₹* ~	>		(a) (b)	} _		
82	\ \(\frac{1}{2}\)	-	V	**************************************	diphosgene (e)	€	₹	

45 Q &	Structura 8	35	30	Synthesis	20	15	10	5
3 3			<u>_</u>				(g) (g)	
82 22	\bigcirc		>	Q O	5 g }			
98				5	, a			
87	D			5	e Hay	On Con N		
98	3		~ <u></u>	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	E, CO,			₹ E

5 10 15 20	Synthesis	HO (ii) (iii) (iv)	(a) (b)	(a) (b)	(a)	C CN LIAIM,	HO NH, NH, (8) (7)	
<i>30</i>	9.17				S HIN	NH ₁		OZ Z
40 45	No Structure	8	s s	£	25	88	\$	88



[0272] The following compounds can be prepared according to the synthesis schemes:



(u) potassium tert. butanolate, THF, 24h, 0 - 50 °C; (v) chromatographic separation; (w) NH $_3$ (fl.), MeOH, -78 - 0 °C.

[0273] Compounds 1 to 75 are prepared according to the following procedures:

METHOD A:

[0274] A solution of 1-bromo-5-phenoxypentane (1.4 to 3.5 mmol) in ten equivalents of the suitable secondary amine was heated to reflux temperature with stirring for 48 hours (compds. 1, 3 and 4), 24 hours (compd. 2) or 4 hours (compd. 5). After cooling, the excess base was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate formed was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD B:

10

[0275] A solution of 1-bromo-5-phenoxypentane (0.9 to 1.7 mmol) and an excess of the suitable secondary amine (2.3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 48 hours (compd. 6) or 24 hours (compds. 7, 8, 9, 10, 11, 12&13, 14, 15, 16, 17 and 29). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 12 and 13 were separated by column chromatography on silica gel eluting with a solvent mixture of petroleum spirit (bp 60-80°C), diethyl ether and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compounds 14 and 16 were purified by column chromatography on silica gel eluting with diethyl ether and triethylamine in the ratio 99:1, and the eluent was removed under reduced pressure to leave an oil. The oil was converted to oxalate salt (compds. 6, 7, 8, 9, 11, 12, 13, 15, 16, 17 and 29) by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from isopropyl alcohol (compds. 6, 7, 10, 13 and 16), absolute ethanol (compds. 8, 9, 11, 12, 15 and 29) or methanol (compd. 17). The oil was converted to hydrochloride salt (compd. 14) by adding 2N HCI. The precipitate was formed in a mixture of chloroform and diethyl ether (1:1) and recrystallised from acetone.

METHOD C:

[0276] A solution of the suitable α -bromo- ω -aryloxy alkane (0.4 to 1.4 mmol) or ω -bromoalkyl phenyl sulphide (1 mmol, compds. 33 and 34) and an excess of pyrrolidine (10 to 15 equivalents) or 3-methylpiperidine (10 equivalents, compd. 38) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours or 16 hours (compd. 47). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD D:

40

[0277] A solution of the suitable 4'-(5-bromopentoxy)phenyl ketone (0.7 to 1 mmol, compds. 39, 44 and 45) or 1-bromo, 5-(4-phenoxyphenoxy)pentane (0.6 mmol, compd. 48) and an excess of pyrrolidine (10 to 15 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours (compds. 39, 44 and 48) or 24 hours (compd. 45). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with chloroform (compds. 39, 45 and 48) or dichloromethane (compd. 44), the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (recrystallised twice from absolute ethanol in the case of compd. 39).

METHOD E:

55 **[0278]**

1. The oxalate 18 was prepared according to method C. A solution of compound 18 (0.57 mmol) in 10 ml methanol and 10 ml absolute ethanol was placed with 100 mg of palladium (5%) on carbon catalyst in a two-neck round-bot-

tom flask fitted with a balloon filled with hydrogen. The mixture was stirred vigorously at room temperature and the flask was purged of air and filled with hydrogen. After 3 hours, the catalyst was filtered off on celite and the solvent removed under reduced pressure. The residual solid was converted to oxalate salt by dissolving in methanol and adding a solution of oxalic acid (2 equivalents) in absolute ethanol. Diethyl ether was added to form a precipitate. The product was recrystallised from absolute ethanol.

2. To a solution of compound 40 (0.35 mmol) in pyridine vigorously stirred at 0°C was added dropwise a slight excess of benzoyl chloride (0.4 mmol). The stirring was allowed to continue 20 minutes after the end of the addition after which the mixture was placed in the refrigerator overnight (16 hours). The solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with chloroform, the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was dissolved in methanol, filtered, and concentrated under reduced pressure, the solid was recrystallised from absolute ethanol

15 METHOD F:

5

10

[0279] In a three-neck flask kept under nitrogen was placed a solution of the suitable phenol (1.6 mmol), 3-(diethylamino)propanol (1.5 mmol), and triphenyl phosphine (1.9 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diisopropyl azodicarboxylate (2 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCI (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (compds. 43 and 46) or from a 1:1 mixture of methanol and absolute ethanol (compd. 42).

30 METHOD G:

[0280] A solution of the free base of compound 39 (0.6 mmol) or compound 46 (0.8 mmol) in 20 ml dry diethyl ether was added dropwise to a stirred suspension of lithium aluminium hydride (0.6 or 0.8 mmol) in 20 ml dry diethyl ether kept under nitrogen. The mixture was stirred at room temperature under nitrogen for two hours. Ice-cold water was carefully added and the organic layer decanted. The aqueous phase was extracted with diethyl ether. The combined organic solutions were dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a yellow oil. The oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (compd 50) or from isopropyl alcohol, giving a very hygroscopic solid (compd. 63).

METHOD H:

40

55

[0281] A solution of the suitable α-bromo-ω-(4-cyanophenoxy) alkane (0.5 to 0.7 mmol) and an excess of the suitable secondary amine (8 to 12 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours (compds. 54, 55, 57 and 60), 20 hours (compd. 52), 16 hours (compds. 56, 58, 59 and 61) or 8 hours (compd. 51) or was stirred at room temperature for 48 hours (compd. 53) or 24 hours (compd. 60). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 62 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. For all the compounds of method H, the remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (two recrystallisations were required for compds. 58 and 59) or from a 1:1 mixture of methanol and absolute ethanol (compd. 55).

METHOD J:

[0282] A solution of compound 46 (1 mmol) in 10 ml methanol was stirred at room temperature and a solution of

hydroxylamine hydrochloride (2 equivalents) in 2 ml water was added. The mixture was stirred at 50-70°C in a water bath for 20 minutes. Methanol was removed under reduced pressure. The residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 64 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. Diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

10 METHOD K:

[0283] A solution of 4'-(3-bromopropoxy)acetophenone (0.8 to 1.9 mmol) and an excess of the suitable piperidine (3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours. After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 67 and 70 were separated by column chromatography on silica gel eluting with a solvent mixture of diethyl ether, petroleum spirits (bp 60-80°C) and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compound 75 was purified by column chromatography on silica gel eluting with chloroform and methanol (1:1), and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD L:

25

40

[0284] In a three-neck flask kept under nitrogen was placed a solution of the suitable 4'-hydroxyphenyl ketone (0.9 to 3 mmol), 3-(1-piperidinyl)propanol (0.9 to 3 mmol), and triphenyl phosphine (1 to 3.5 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diethyl azodicarboxylate (1 to 3.6 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCl (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The crude product was purified by column chromatography on silica gel eluting with diethyl ether containing 1% triethylamine, and concentrated under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

[0285] Compounds 76 to 97 are prepared according to the following procedures:

Example 76

3,3-Dimethylbutyl 3-piperidinopropyl ether

[0286] Sodium 3-piperidinopropanolate (5 mmol), 5 mmol of 3,3-dimethylbutyl chloride, a catalytic amount of tetrabutylammonium iodide, and 0.5 mmol of 15-crown-5 in 10 ml of dry dimethyl sulfoxide were refluxed for 12 hours. Water was added, and it was extracted with diethyl ether. The organic layer was purified by column chromatography on silica gel (eluent: methylene chloride/methanol (90/10), ammonia atmosphere). The solvent was removed under reduced pressure and the residue crystallized with oxalic acid from diethyl ether/ethanol.

SF: $C_{14}H_{29}NO \times 1.1 C_2H_2O_4$ (326.4)

mp: 143 °C

55

CHN analysis	calculated:	C 59.6	H 9.63	N 4.29
	found:	C 59.7	H 9.61	N 4.30

Example 77

3-Phenylpropyl 3-piperidinopropyl ether

[0287] Sodium 3-piperidinopropanolate (20 mmol), 20 mmol of 3-phenylpropyl bromide, and 0.5 mmol of 15-crown-5 in 30 ml of dry toluene were refluxed for 4 hours. The solvent was evaporated and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol/aqueous ammonia (90/10/0.5)). After removing the solvent under reduced pressure the residue was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₇H₂₇NO x C₂H₂O₄ (351.4)

mp: 125 °C

CHN analysis calculated: C 64.9 H 8.32 N 3.99 found: C 64.9 H 8.13 N 4.02

Example 78

3-(4-Chlorophenyl)propyl 3-piperidinopropyl ether

[0288] Sodium 3-piperidinopropanolate (20 mmol), 7 mmol of 3-(4-chlorophenyl)propylmesylate, and 0.5 mmol of 15-crown-5 in 30 ml of dry toluene were refluxed for 4 hours. The solvent was evaporated and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol (90/10)). After removing the solvent under reduced pressure the residue was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₇H₂₆NOCl x C₂H₂O₄ (385.9)

mp: 147 °C

30

10

15

CHN analysis	calculated:	C 59.1	H 7.31	N 3.63
	found:	C 59.0	H 7.34	N 3.60

35 Example 79

2-Benzothiazolyl 3-piperidinopropyl ether

[0289] Sodium 3-piperidinopropanolate (5 mmol) and 5 mmol of 2-chlorobenzothiazole in
 20 ml of dry tetrahydrofurane were refluxed for 12 hours. The suspension was filtered and the solvent evaporated under reduced pressure. The product was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₅H₂₀N₂OS x C₂H₂O₄ (366.4)

mp: 178.2-178.8 °C

45

50

CHN analysis	calculated:	C 55.7	H 6.05	N 7.64
	found:	C 55.6	H 6.03	N 7.51

Example 80

N-Phenyl-3-piperidinopropyl carbamate

[0290] 3-Piperidinopropanol hydrochloride (10 mmol) and 10 mmol of phenyl isocyanate in 40 ml of dry acetonitrile were refluxed for 3 hours. The solvent was evaporated, and then the residue was recrystallized in dry ethanol.

SF: C₁₅H₂₂N₂O₂ x HCl x 0.1 H₂O (300.6)

mp: 169-170 °C

 CHN analysis
 calculated:
 C 59.9
 H 7.78
 N 9.32

 found:
 C 59.9
 H 7.64
 N 9.05

Example 81

5

20

35

40

55

10 N-Pentyl-3-piperidinopropyl carbamate

[0291] 3-Piperidinopropanol hydrochloride (4 mmol) and 4 mmol of pentyl isocyanate in 20 ml of dry acetonitrile were refluxed for 3 hours. The solvent was evaporated and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol/aqueous ammonia (90/10/0.5)). After removing the solvent under reduced pressure the residue was crystallized with hydrochloric acid in 2-propanol.

SF: C₁₄H₂₈N₂O₂ x HCl x 0.5 H₂O (301.9) mp: 88-89 °C

CHN analysis calculated: C 55.7 H 10.0 N 9.28 found: C 55.7 H 9.84 N 9.18

25 Example 82

(S)-(+)-N-[2-(3,3-Dimethyl)butyl]-3-piperidinopropyl carbamate

[0292] 3-Piperidinopropanol hydrochloride (5 mmol) and 5 mmol of (S)-2-(3,3-dimethyl)butyl isocyanate in 10 ml of dry acetonitrile were refluxed for 12 hours. The solvent was evaporated and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol (90/10), ammonia atmosphere). The solvent was removed and the residue crystallized with oxalic acid from diethyl ether/ethanol.

SF: $C_{15}H_{30}N_2O_2 \times C_2H_2O_4 \times 0.25 H_2O$ (365.0) mp: 148 °C [α]²³ = +10.4° (c = 0.495, Methanol)

 CHN analysis
 calculated:
 C 56.0
 H 8.98
 N 7.68

 found:
 C 56.0
 H 9.01
 N 7.64

Example 83

45 N-(4-Chlorobenzyl)-S-(3-piperidinopropyl) isothiourea

[0293] 4-Chlorobenzylamine (10 mmol) was added dropwise to 10 mmol of benzoylisothiocyanate dissolved in 20 ml of dry ether followed by stirring for 2 hours. The precipitated product was filtered off and crystallized from ethyl acetate (Yield: 60%). Potassium carbonate (10 mmol) in 30 ml of water was added dropwise to 5 mmol of the product in 20 ml of ethanol and refluxed for 2 hours. The precipitated product was filtered off and crystallized from ethyl acetate/petroleum ether (Yield: 65%). 3-Piperidinopropyl chloride hydrochloride (3 mmol), 3 mmol of the product, and a catalytic amount of potassium iodide were refluxed in 20 ml of ethanol for 2 days. Subsequently the ethanol was evaporated and the residue purified by column chromatography using methanol/ethyl acetate (2/8) as eluent. After evaporation of the solvent, the product was crystallized with hydrochloric acid from diethyl ether/ethanol.

SF: C₁₆H₂₄CIN₃S x 2 HCl x H₂O (416.8) mp: 104-107.5 °C

 CHN analysis
 calculated:
 C 46.1
 H 6.77
 N 10.1

 found:
 C 45.9
 H 6.87
 N 9.69

Example 84

5

15

20

25

N'-Cyclohexylthiocarbamoyi-N-1,4'-bipiperidine

[0294] 1,4'-Bipiperidine (5 mmol) in 10 ml of dry ether was added dropwise to 5 mmol of cyclohexyl isothiocyanate in 30 ml of dry ether followed by stirring for 2 hours. Filtration gave a residue, which was dissolved in ethanol and crystallized with oxalic acid. Recrystallization resulted in the pure product.

SF: C₁₇H₃₁N₃S x H₂C₂O₄ x 0.25 H₂O (404.1)

mp: 225-226 °C

CHN analysis	calculated:	C 56.5	H 8.35	N 10.39
	found:	C 56.2	H 8.25	N 10.33

Example 85

N-Heptanoyl-1,4'-bipiperidine

[0295] 1,4'-Bipiperidine (10 mmol) in 5 ml of water was added dropwise to a solution of 5 mmol of n-heptanoyl chloride in 20 ml of dioxane. After stirring for 15 minutes the solvent was evaporated under reduced pressure and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol/aqueous ammonia (90/10/0.5)). The solvent was removed under reduced pressure, and the residue was crystallized with oxalic acid.

SF: C₁₇H₃₂N₂O x H₂C₂O₄ (370.5)

mp: 131-132 °C

35

40

CHN analy	/sis	calculated:	C 61.6	H 9.25	N 7.56
		found:	C 61.6	H 9.36	N 7.50

Example 86

3-Cyclopentyl-N-(3-(1-pyrrolidinyl)propyl)propanamide

[0296] 3-Cyclopentyl propionylchloride (5 mmol) in 10 ml of dioxane was added dropwise to a solution of 10 mmol of 1-(3-aminopropyl)pyrrolidine in water. After stirring for 4 hours the solvent was evaporated under reduced pressure and the residue purified by column chromatography on silica gel (eluent: methylene chloride/ methanol/aqueous ammonia (90/10/1)). The solvent was removed under reduced pressure and the residue was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₇H₂₈N₂O x H₂C₂O₄ x 0.5 H₂O (351.2)

mp: 89.5 °C

55

CHN analysis	calculated:	C 58.1	H 8.83	N 7.97
	found:	C 58.1	H 8.76	N 7.87

Example 87

N-Cyclohexyl-N'-(1-pyrrolidinyl-3-propyl)urea

[0297] In an argon atmosphere 10 mmol of cyclohexylisocyanate was added slowly to 10 mmol of 1-(3-aminopro-pyl)pyrrolidine in 10 ml of acetonitrile. The product preci-pitated instantly as a pure white solid. The solvent was removed under reduced pressure and the product was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₄H₂₇N₃O x C₂H₂O₄ x 0.25 H₂O (347.7) Yield: 83%

mp: 113.3 °C

10

15

CHN analysis	calculated:	C 56.0	H 8.45	N 12.2
	found:	C 55.6	H 8.27	N 12.0

Example 88

α-(4-Acetylphenoxy)-α'-piperidino p-xylol

20

[0298] Hydroxyacetophenone (2 mmol) and 5 mmol of K_2CO_3 were stirred in 20 ml of acetone with 2 ml of DMF for 10 minutes. After addition of 3.5 mmol of α,α' -dibromoxylol the reaction was stirred at ambient temperature for 12 hours and after addition of 7 mmol of piperidine for 1 hour under reflux. The solvent was evaporated under reduced pressure. The residue was suspended in water, extracted with methylene chloride. The combined organic extracts were crystallized with oxalic acid. Recrystallization resulted in the pure product.

SF: C₂₁H₂₅NO₂ x C₂H₂O₄ (413.5)

mp: 136-137 °C

30

CHN analysis	calculated:	C 66.8	H 6.58	N 3.39
	found:	C 66.7	H 6.70	N 3.40

35 Example 89

α -(4-Acetylphenoxy)- α '-(1-pyrrolidinyl) p-xylol

[0299] Hydroxyacetophenone (2 mmol) and 5 mmol of K_2CO_3 were stirred in 20 ml of acetone with 2 ml of DMF for 10 minutes. After addition of 3.5 mmol of α,α' -dibromoxylol the reaction was stirred at ambient temperature for 12 hours and after addition of 7 mmol of pyrrolidine for 1 hour under reflux. The solvent was evaporated under reduced pressure. The residue was suspended in water, extracted with methylene chloride. The combined organic extracts were crystallized with oxalic acid. Recrystallization resulted in the pure product.

SF: C₂₀H₂₃NO₂ x C₂H₂O₄ x 0.25 H₂O (404.0)

mp: 136-137 °C

50

CHN analysis	calculated:	C 65.4	H 6.36	N 3.47
	found:	C 65.6	H 6.29	N 3.47

Example 90

α -(3-Phenylpropoxy)- α '-piperidino p-xylol

[0300] 4-(Piperidinomethyl)benzoic acid methyl ester (22 mmol) in dry tetrahydrofurane was added dropwise to a suspension of 44 mmol of lithium aluminium hydride in 30 ml of dry tetrahydrofurane at 0 °C. After refluxing for 2 hours a

saturated solution of ammonium chloride in water was added dropwise. After stirring for 12 hours at ambient temperature the organic layer was isolated and the aqueous layer extracted with methylene chloride. The organic extracts were combined and the solvent was evaporated under reduced pressure. The residue was crystallized with maleic acid from diethyl ether/2-propanol (Yield: 91%). Sodium 4-(piperidinomethyl)benzyl alcoholate (5 mmol) and 6 mmol of 3-phenyl-propyl bromide in 10 ml of dry toluene were refluxed for 6 hours. The solvent was evaporated under reduced pressure. The residue was purified by rotatory chromatography on silica gel using methylene chloride/ammonia atmosphere as eluent. The product was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₂₂H₂₉NO x C₂H₂O₄ x 0.5 H₂O (422.5)

mp: 104-105 °C

10

15

CHN analysis	calculated:	C 68.2	H 7.63	N 3.32
	found:	C 68.3	H 7.26	N 3.36

Example 91

3-(4-Chlorobenzyl)-5-(2-piperidinoethyl)-1,2,4-oxadiazole

20

[0301] Hydroxylamine hydrochloride (20 mmol) was added dropwise to a solution of 20 mmol of sodium in 50 ml of methanol at 0 °C. After stirring for 30 minutes at ambient temperature 10 mmol of 4-chlorobenzyl cyanide was added dropwise at

0 °C. After refluxing for 6 hours the suspension was filtered and the solvent evaporated under reduced pressure. The residue was crystallized from diethyl ether (Yield: 41%). To a solution of 4 mmol of the product and 6 mmol of 3-piperidinopropionic acid methyl ester in 15 ml of dry methanol 5 mmol of sodium in 20 ml of methanol was added dropwise at 0 °C. After stirring for 1 hour under argon atmosphere followed by refluxing for 18 hours the solvent was evaporated under reduced pressure. The residue was suspended in DMF and stirred for 6 hours at 80 °C. The solvent was evaporated under reduced pressure. The residue was suspended in water and extracted with methylene chloride. The residue of the organic layer was purified by rotatory chromatography on silica gel using methylene chloride/ammonia atmosphere as eluent. The product was crystallized with oxalic acid from diethyl ether/ethanol.

SF: $C_{16}H_{20}CIN_3O \times C_2H_2O_4$ (395.8)

mp: 152-154 °C

35

40

CHN analysis	calculated:	C 54.6	H 5.60	N 10.6
	found:	C 54.3	H 5.60	N 10.5

_

Example 92

2-((2-Piperidinoethyl)amino)benzothiazole

[0302] 2-Chlorobenzothiazole (10 mmol), 10 mmol of 2-piperidinoethanamine, and 30 mmol of triethylamine in 50 ml of dry ethanol were refluxed for 6 hours. The product was crystallized with hydrochloric acid in 2-propanol and recrystallized in methanol.

SF: C₁₄H₁₉N₃S x 2 HCl x 0.25 H₂O (338.8)

Yield: 95%

mp: 225 °C

55

CHN analysis	calculated:	C 49.6	H 6.40	N 12.4
	found:	C 49.5	H 6.49	N 12.3

Example 93

5-Piperidinopentylamine

[0303] 5-Chlorovaleronitrile (10 mmol), 20 mmol of piperidine, 20 mmol of potassium carbonate and a catalytic amount of potassium iodide in 50 ml of ethanol were refluxed for 6 hours. The solvent was removed under reduced pressure, the residue suspended in water and extracted with methylene chloride. The organic layer was purified by column chromatography on silica gel using methylene chloride/methanol/aqueous ammonia (90/10/1) as eluent (Yield: 59%). The product was added dropwise to a suspension of 25 mmol of lithium aluminium hydride in 25 ml of dry tetrahydrofurane at 0 °C. After refluxing for 1 hour 10 ml of a saturated solution of sodium/potassium tartrate in water was added dropwise. The residue was filtered off and the filtrate purified by column chromatography on silica gel using methylene chloride/methanol/aqueous ammonia (90/10/1) as eluent. The residue was crystallized with hydrochloric acid from diethyl ether/2-propanol.

SF: C₁₀H₂₂N₂ x 2 HCl x 0.5 H₂O (252.2)

mp: 187 °C

20

15

CHN analysis	calculated:	C 47.6	H 9.99	N 11.1
	found:	C 47.8	H 9.70	N 11.0

Example 94

5 5-Nitro-2-(6-piperidinohexyl)pyridine

[0304] 6-Aminohexanol (15 mmol), 15 mmol of 2-chloro-5-nitropyridine, 5 ml of triethylamine, and a catalytic amount of potassium iodide were refluxed in 30 ml of ethanol for 12 hours. The solvent was evaporated, and the residue was purified by column chromatography on silica gel (eluent: methylene chloride/methanol (95/5), ammonia atmosphere). The solvent was removed under reduced pressure (Yield: 66%). The product (5 mmol) was dissolved in tetrahydrofurane, stirred at 0 °C and 10 mmol of thionyl chloride was added dropwise. After 1 hour at ambient temperature the mixture was warmed to 60 °C for 2 hours. The solvent and the excess of thionyl chloride were evaporated. The oily residue was crystallized with hydrochloric acid from diethyl ether/ethanol (Yield: 95%). The product (5 mmol), 10 mmol of piperidine, 15 mmol of potassium carbonate, and a catalytic amount of potassium iodide were refluxed in 30 ml of ethanol for 12 hours. The solvent was evaporated and the residue purified by column chromatography (eluent: methylene chloride/methanol (95/5), ammonia atmosphere). The solvent was removed under reduced pressure, and the residue was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₆H₂₆N₄O₂ x C₂H₂O₄ (396.4)

mp: 118.6-119.7 °C

45

40

CHN analysis	calculated:	C 54.5	H 7.12	N 14.1
	found:	C 54.4	H 7.18	N 14.2

Example 95

3-Nitro-2-(6-piperidinohexylamino)pyridine

[0305] 6-Aminohexanol (15 mmol), 15 mmol of 2-chloro-3-nitropyridine, 5 ml of triethylamine and a catalytic amount of potassium iodide were refluxed in 30 ml of ethanol for 12 hours. The solvent was evaporated and the residue was purified by column chromatography on silica gel (eluent: methylene chloride/methanol (98/2), ammonia atmosphere). The solvent was removed under reduced pressure (Yield: 55%). The product (5 mmol) was dissolved in tetrahydrofurane, stirred at 0 °C and 10 mmol of thionyl chloride was added dropwise. After 1 hour at ambient temperature the mixture was warmed to 60 °C for 2 hours. The solvent and the excess of thionyl chloride were evaporated. The oily residue was crystallized with hydrochloric acid from diethyl ether/ethanol (Yield: 95%). The product (5 mmol), 10 mmol of piperidine, 15 mmol of potassium carbonate, and a catalytic amount of potassium iodide were refluxed in 30 ml of eth-

anol for 12 hours. The solvent was evaporated and the residue purified by column chromatography (eluent: methylene chloride/methanol (95/5), ammonia atmosphere). The solvent was removed under reduced pressure, and the residue was crystallized with oxalic acid from diethyl ether/ethanol

SF: C₁₆H₂₆N₄O₂ x C₂H₂O₄ (396.4)

mp: 130.3-130.7 °C

10

5

CHN analysis	calculated:	C 54.5	H 7.12	N 14.1
	found:	C 54.3	H 7.14	N 13.9

Example 96

5 2-(6-Piperidinohexylamino)pyrimidine

[0306] 6-Aminohexanol (15 mmol), 15 mmol of 2-chloropyrimidine, 5 ml of triethylamine, and a catalytic amount of potassium iodide were refluxed in 30 ml of ethanol for 12 hours. The solvent was evaporated, and the residue was purified by column chromatography on silica gel (eluent: methylene chloride/methanol (98/2), ammonia atmosphere). The solvent was removed under reduced pressure (Yield: 40%). The product (5 mmol) was dissolved in tetrahydrofurane, stirred at 0 °C and 10 mmol of thionyl chloride was added dropwise. After 1 hour at ambient temperature the mixture was warmed to 60 °C for 2 hours. The solvent and the excess of thionyl chloride were evaporated. The oily residue was crystallized with hydrochloric acid from diethyl ether/ethanol (Yield: 95%). The product (5 mmol), 10 mmol of piperidine, 15 mmol of potassium carbonate, and a catalytic amount of potassium iodide were refluxed in 30 ml of ethanol for 12 hours. The solvent was evaporated and the residue purified by column chromatography (eluent: methylene chloride/methanol (95/5), ammonia atmosphere). The solvent was removed under reduced pressure, and the residue was crystallized with oxalic acid from diethyl ether/ethanol.

SF: C₁₅H₂₆N₄ x C₂H₂O₄ (352.4)

mp: 150.3-150.9 °C

30

35

CHN analysis	calculated:	C 57.9	H 8.00	N 15.9
	found:	C 58.0	H 8.14	N 15.8

Example 97

N-(6-Phenylhexyl)piperidine

40

[0307] 6-Phenylhexanol (5 mmol) was stirred at 0 °C, and thionyl chloride (10 mmol) was added dropwise. After 1 hour at ambient temp. the mixture was warmed to 60 °C for 2 hours. The excess of thionyl chloride was evaporated. The oily residue was purified by column chromatography on silica gel (eluent: methylene chloride) (Yield: 98%). The product was dissolved in 50 ml of ethanol, and 10 mmol of K_2CO_3 , 1 mmol of Kl, and 10 mmol of piperidine were added. After refluxing for 6 hours the solvent was evaporated under reduced pressure. The residue was suspended in water and extracted with methylene chloride. The organic extracts were combined, dried with MgSO₄ and the residue purified by column chromatography on silica gel (eluent: methylene chloride/methanol/aqueous ammonia (90/10/1)). The residue was crystallized with oxalic acid from diethyl ether/methanol.

SF: C₁₇H₂₇N x C₂H₂O₄ (335.5)

mp: 152 °C

55

CHN analysis	calculated:	C 68.0	H 8.71	N 4.18
	found:	C 68.0	H 8.67	N 4.05

Pharmacological study

[0308] Interaction of compounds with the H_3 receptor are evidenced *in vitro* by the measurement of the release of neosynthesized tritiated histamine from rat cerebral cortex synaptosomes preincubated with tritiated histidine (Garbarg et al., J. Pharmacol. Exp. Ther., 1992, 263: 304-310). The H_3 potency of agonists is measured by the inhibition of tritiated histamine release and that of antagonists by the progressive reversal of release inhibition by the selective H_3 agonist (R) α -methylhistamine (Arrang et al., Nature, 1987, 327: 117-123).

[0309] Interaction of compounds with the H_3 receptor are evidenced in vitro on guinea-pig ileum by the procedure described by Ligneau et al., J. Pharmacol. Exp. Ther. 271, 452-459 (1994).

[0310] Briefly, longitudinal muscle strips from guinea-pig small intestine were dissected out and incubated in a gassed O₂/CO₂ (95 %/5 %) modified Krebs-Ringer's bicarbonate medium at +37°C in presence of 1 μM mepyramine to block the H₁ receptor. After equilibration, contractile activity under stimulation (rectangular pulses of 15 V, 0,5 msec, 0,1 Hz) was recorded.

[0311] Concentration-response curves of the effect of $(R)\alpha$ -Methylhistamine alone or together with the antagonist were established.

[0312] The effects of agonists and antagonists were estimated *in vivo* by the measurement of the tele-methylhistamine level variations in the brain of mice (Garbarg et al., J. Neurochem., 1989, 53: 1724-1730). At various time after p.o. administration of the compounds, the effect of agonists and antagonists are evidenced by the decrease and increase respectively in telemethylhistamine level induced.

[0313] The changes are compared to those induced by reference compounds given in high dosage and this allows the calculation of the ED₅₀ value for each compound which correspond to the dose responsible for an half maximal effect.

[0314] The results are reported in the following tables II and III:

25

TABLE II

30
35
40

Ex No.	X	n	R ¹ R ²	$R^3 (n_3 = 1)$	Ki (nM)	ED ₅₀ (mg/kg/p.o.)	
18	0	5	-(CH ₂) ₄ -	p-NO ₂	39 ± 11	1.1	
43	0	3	Et, Et	p-CN	95 ± 28	0.50	
46	0	3	Et, Et	p-CH ₃ CO		0.44	
50	0	5	-(CH ₂) ₄ -	p-CH ₃ CH(OH)		1.0	
56	0	4	Et, Et	p-CN		1.1	
59	0	3	-(CH ₂) ₅ -	p-CN		0.20	
60	0	3	-(CH ₂) ₆ -	p-CN		0.64	
63	0	3	Et, Et	p-CH ₃ CH(OH)		0.34	
64	0	3	Et, Et	p-CH ₃ C=N(OH)		0.8	
66	0	3	-(3-Me)-(CH ₂) ₅ -	p-CH ₃ CO	0.3		
68	0	3	-(4-Me)-(CH ₂) ₅ -	p-CH ₃ CO		0.3	
69	0	3	-(CH ₂) ₅ -	p-C ₂ H ₅ CO		0.4	

50

45

H₃-receptor antagonist activity pA₂ (guineaplig ileum)

81 6.3

85 6.4

91 7.2

TABLE III

TABLE III (continued)

Example No.	H ₃ -receptor antagonist activity pA ₂ (guineaplg ileum)
92	6.6
97	6.5

[0315] All the above compounds 1 to 97 were find to be H_3 -antagonists.

[0316] Comparative data concerning the activity of imidazole derivatives and of the non-imidazole analogues according to the invention, are reported below in Table IV:

TABLE IV:

5	Imidazole derivative	Non-imidazole analogue according to the invention
10	$\begin{array}{c} \\ \text{HN} \\ \\ N \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \end{array} = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$	ex 59: NC————————————————————————————————————
15	Ki = 12 nM $ED_{50} = 0.54 \text{ mg/kg}$	Ki = 11 nM ED ₅₀ = 0,20 mg/kg ex 43:
20		NC-CH ₂ CH ₂ CH ₃ CH ₂ CH ₃
25		Ki = 95 nM ED ₅₀ = 0.50 mg/kg ex 58:
30		NC-(CH ₂) ₃ -N
35		Ki = 20 nM ex 60:
40		NC $ O-(CH_2)_3$ $ N$
45	(CH ₂) ₃ -O-(CH ₂) ₃ -O	ex 77:
50	HN N Ki = 17 nM	Ki = 15 mM

Claims

5

15

20

30

35

40

50

55

1. Compound of general formula (A):

 $[W] - N \stackrel{R^1}{\underset{R^2}{\longleftarrow}} (A)$

in which:

- W is a residue which imparts antagonistic and/or agonistic activity at histamine H₃-receptors when attached to an imidazole ring in 4(5)-position;
- R¹ and R² may be identical or different and represent each independently
 - a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
 - · a saturated nitrogen-containing ring

i) N (CR^aR^b)_m

with m ranging from 2 to 8, or

a non-aromatic unsaturated nitrogen-containing ring

(CHRa)_p CRb

with p and q being from 0 to 3 independently and r being from 0 to 4, provided that p and q are not simulteously 0 and $2 \le p + q + r \le 8$,

Ra-d being independently a hydrogen atom or a lower alkyl, cycloalkyl, or carboalkoxy group, or

- · a morpholino group, or
- a N-substituted piperazino group:

45 -N-R

with R being a lower alkyl, cycloalkyl, carboalkoxy, aryl, arylalkyl, an alkanoyl or aroyl group, as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers.

- 2. Compound according to claim 1, in which R¹ and R² are independently a lower alkyl group.
- 3. Compound according to claim 2, in which R¹ and R² are each an ethyl group.
- 4. Compound according to claim 1, in which —NR¹R² is a saturated nitrogen-containing ring:

i) N (CHR^a)_m

m being as defined in claim 1.


5

15

20


25

- 5. Compound according to claim 4, characterized in that m is 4, 5 or 6.
 - 6. Compound according to claim 5, characterized in that -NR¹R² represents a piperidyl group.
 - 7. Compound according to claim 5, characterized in that -NR¹R² represents a pyrrolidinyl group.
 - 8. Compound according to claim 1, characterized in that —NR¹R² is a non-aromatic unsaturated nitrogen-containing ring:



Ra-d and p, q and r being as defined in claim 1.

- 9. Compound according to claim 8, characterized in that p, q and r are 1 or 2, more preferably p is 2 and q and r are 1.
 - 10. Compound according to anyone of claims 4 to 9, characterized in that R^{a-d} represents each an hydrogen atom.
- 11. Compound according to anyone of claim 4 to 9, characterized in that the nitrogen-containing ring i) or ii) is substituted, preferably mono- or di-substituted, more preferably mono-substituted, with an alkyl group.
 - 12. Compound according to claim 11, characterized in that the nitrogen-containing ring is mono-substituted with a methyl group.
- 40 13. Compound according to anyone of claims 11 and 12, characterized in that the substituent(s) is(are) in meta-position with respect to the nitrogen atom.
 - 14. Compound according to claim 1, characterized in that —NR¹R² is a morpholino group.
- 45 15. Compound according to claim 1, characterized in that —NR¹R² is a N-substituted piperazino group, preferably N-acetylpiperazino.
 - 16. Compound according to anyone of claims 1 to 15, of general formula (I):



in which:

 C_nH₂n is a linear or branched hydrocarbon chain with n ranging from 2 to 8; — X is an oxygen or sulfur atom; R¹ and R² are as defined in claim 1; — n₃ is an integer from 0 to 5. 5 17. Compound according to claim 16, characterized in that n₃ is zero. 18. Compound according to anyone of claims 16 and 17, characterized in that n₃ is 1 with R³ being as defined in claim 1 and preferably in para-position. 10 19. Compound according to anyone of claims 16 and 18, characterized in that R³ is a lower alkyl, preferably a C₁-C₄ alkyl. 20. Compound according to anyone of claims 16 and 18, characterized in that R³ is a halogen atom, a cyano, nitro, alkanoyl, alkyloximine or hydroxyalkyl, preferably CN, NO2, COCH3, COC2H5, H3C-C=N-OH or H3C-CHOH. 15 21. Compound according to claim 16, characterized in that R3 taken together with the carbon atoms of the phenyl group to which it is fused, form a 5- or 6- membered saturated or unsaturated ring, in particular a 5,6,7,8-tetrahydronaphthyl group. 20 22. Compound according to claim 16, characterized in that R³ taken together with the phenyl group to which it is fused, form a naphthyl group. 23. Compound according to anyone of claims 16 to 22, characterized in that -C_nH_{2n}- is a linear hydrocarbon chain -(CH₂)_n-, n being as defined in claim 16. 25 24. Compound according to anyone of claims 16 to 23, characterized in that X is an oxygen atom. 25. Compound according to anyone of claims 16 to 23, characterized in that X is a sulfur atom. 30 26. Compound according to anyone of claims 16 to 25, characterized in that n is varying from 3 to 5 and is preferably 3. 27. Compound according to anyone of claims 16 to 26, characterized in that it is one of the following compounds: 1-(5-phenoxypentyl)-piperidine 35 1-(5-phenoxypentyl)-pyrrolidine N-methyl-N-(5-phenoxypentyl)-ethylamine 1-(5-phenoxypentyl)-morpholine N-(5-phenoxypentyl)-hexamethyleneimine N-ethyl-N-(5-phenoxypentyl)-propylamine 40 1-(5-phenoxypentyl)-2-methyl-piperidine 1-(5-phenoxypentyl)-4-propyl-piperidine 1-(5-phenoxypentyl)-4-methyl-piperidine 1-(5-phenoxypentyl)-3-methyl-piperidine 1-acetyl-4-(5-phenoxypentyl)-piperazine 45 1-(5-phenoxypentyl)-3,5-trans-dimethyl-piperidine 1-(5-phenoxypentyl)-3,5-cis-dimethyl-piperidine 1-(5-phenoxypentyl)-2,6-cis-dimethyl-piperidine 4-carboethoxy-1-(5-phenoxypentyl)-piperidine 3-carboethoxy-1-(5-phenoxypentyl)-piperidine 50 1-(5-phenoxypentyl)-1,2,3,6-tetrahydropyridine 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine 1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine 1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine 1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine 55 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine

1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine 1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine

```
1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
             1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
             1-{5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
             1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
             1-(5-phenoxypentyl)-2,5-dihydropyrrole
5
             1-{5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
             1-(4-phenoxybutyl)-pyrrolidine
             1-(6-phenoxyhexyl)-pyrrolidine
             1-(5-phenylthiopentyl)-pyrrolidine
             1-(4-phenylthiobutyl)-pyrrolidine
10
             1-(3-phenoxypropyl)-pyrrolidine
             1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
             1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
             1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
             1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
15
             1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
             1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
             N-[3-(4-nitrophenoxy)-propyl]-diethylamine
             N-[3-(4-cyanophenoxy)-propyl]-diethylamine
             1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
20
             1-{5-[4-(phenylacetyl)-phenoxy]-pentyl}-pyrrolidine
             N-[3-(4-acetylphenoxy)-propyl]-diethylamine
             1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
             1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
             1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
25
             1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
             1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
             1-[5-(4-cyanophenoxy)-pentyl]-piperidine
             N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
             N-[2-(4-cyanophenoxy)-ethyl]-diethylamine
30
             N-[3-(4-cyanophenoxy)-propyl]-dimethylamine
             N-[4-(4-cyanophenoxy)-butyl]-diethylamine
             N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
             1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
              1-[3-(4-cyanophenoxy)-propyl]-piperidine
35
             N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
             N-[6-(4-cyanophenory)-hexyl]-diethylamine
             N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
             N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
             4-(3-diethylaminopropoxy)-acetophenone-oxime
40
              1-[3-(4-acetylphenoxy)-propyl]-piperidine
              1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
              1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine
              1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
              1-[3-(4-propionylphenoxy)-propyl]-piperidine
45
              1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
              1-[3-(4-formylphenoxy)-propyl]-piperidine
              1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
              N-[3-(4-propionylphenoxy)-propyl]-diethylamine
              1-[3-(4-buryrylphenoxy)-propyl]-piperidine
50
              1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine.
     28. Compound according to anyone of claims 16 to 27, characterized in that it is one of the following compounds:
              1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
55
              1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
              1-[3-(4-cyanophenoxy)-propyl]-piperidine
              N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
```

N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine

4-(3-diethylaminopropoxy)-acetophenone-oxime

1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine

1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine

1-[3-(4-propionylphenoxy)-propyl]-piperidine.

29. Compound according to anyone of claims 1 to 15, having the following general formula (IIa) and (IIb):

10
$$\mathbb{R}^1$$
 (chain A ll) — X^1 (chain B ll) — Y^{\parallel} (lla)

or

$$\begin{array}{c|c}
R^{1} \longrightarrow N \longrightarrow (\text{chain A}^{II}) \longrightarrow X^{II} \longrightarrow Y^{II} \\
R^{2} \longrightarrow N \longrightarrow (\text{chain A}^{II}) \longrightarrow X^{II} \longrightarrow Y^{II}
\end{array}$$
(IIb)

20 in which

5

15

25

30

35

45

50

55

- R¹ and R² are as defined with reference to general formula (A) in claim 1;
- the chain A^{II} represents a saturated or unsaturated, straight or branched hydrocarbon chain containing 1 to 6 carbon atoms, it being possible for the saturated hydrocarbon chain to be interrupted by a hetero atom such as a sulphur atom:
- X^{II} represents an oxygen or sulphur atom, -NH-, -NHCO-, -N(alkyl)CO-, -NHCONH-, -NH-CS-NH-, -NHCS-, -O-CO-, -CO-O-, -OCONH-, -OCON(alkyl)-, -OCONH-CO-, -CONH-, -CON(alkyl)-, -SO-, -CO-, -CHOH- or —NR_{II}-C(=NR"_{II})-NR'_{II}-, R_{II} and R'_{II} denoting a hydrogen atom or a lower alkyl radical and R"_{II} a hydrogen atom or another powerful electronegative group, such as a cyano or COY₁ group, Y₁ denoting an alkoxy group;
- the chain B^{II} represents a straight alkylene chain -(CH₂)_{nII}-, n being an integer which can vary between 1 and 5 or a branched alkylene chain containing from 2 to 8 carbon atoms, the alkylene chain being optionally interrupted by one or a number of oxygen or sulphur atoms, or a group -(CH₂)_{nII}-O- or -(CH₂)_{nII}-S- where n_{II} is an integer equal to 1 or 2;
- Y^{II} represents a straight or branched alkyl group containing 1 to 8 carbon atoms; a cycloalkyl containing 3 to 6 carbon atoms; a bicycloalkyl group; a cycloalkenyl group; an aryl group such as an optionally substituted phenyl group; a 5- or 6-membered heterocyclic radical containing one or two heteroatoms chosen from nitrogen and sulphur atoms, the said heterocyclic radical optionally being substituted; or also a bicyclic radical resulting from the fusion of a benzene ring to a heterocycle as defined above.
- 40 30. Compound according to anyone of claims 1 to 15, having the following formula (IIa) and (IIb):

$$\begin{array}{c} R_1 \longrightarrow N \longrightarrow (\text{chain A}^{[l]}) \longrightarrow X^{[l]} \longrightarrow (\text{chain B}^{[l]}) \longrightarrow Y^{[l]} \end{array}$$
 (||a|)

or

$$\begin{array}{c|c} R^1 \longrightarrow N \longrightarrow (\operatorname{chain} A^{[l]}) \longrightarrow X^{[l]} Y^{[l]} \end{array}$$
 (IIb)

in which:

- R¹ and R² are as defined with reference to general formula (A) in claim 1;
- the chain A" represents an unbranched, branched or unsaturated alkyl group -(CH₂)_{nll}- where n_{il} is an integer which can vary between 1 and 8 and preferably between 1 and 4; an unbranched or branched alkene group comprising from 1 to 8 carbon atoms and preferably 1 to 4 carbon atoms; an unbranched or branched alkyne group comprising from 1 to 4 carbon atoms;
- the group XII represents -OCONH-; -OCON(alkyl)-; -OCON(alkene)-; -OCO-; -OCSNH-; -CH₂-; -O-; -

OCH₂CO-; -S-; -CO-; -CS-; amine; alkene;

5

10

15

20

25

30

35

40

45

50

- the chain B^{II} represents an unbranched, branched or unsaturated lower alkyl comprising from 1 to 8 carbon atoms and preferably 1 to 5 carbon atoms; -(CH₂)_{nII}(hetero atom)- where the hetero atom is preferably a sulphur or oxygen atom; n_{II} being an integer which can vary between 1 and 5, preferably between 1 and 4;
- the group YII represents a phenyl group, unsubstituted or mono- or polysubstituted with one or more identical or different substituents selected from halogen atoms, OCF₃, CHO, CF₃, SO₂N(alkyl)₂ such as SO₂N(CH₃)₂, NO2, S(alkyl), S(aryl), SCH2(phenyl), an unbranched or branched alkene, an unbranched or branched alkyne optionally substituted with a trialkylsilyl radical, -O(alkyl), -O(aryl), -CH2CN, a ketone, an aldehyde, a sulphone, an acetal, an alcohol, a lower alkyl, -CH=CH-CHO, -C(alkyl)=N-OH, -C(alkyl)=N-O(alkyl) and other keto derivatives, -CH=NOH, -CH=NO(alkyl), and other aldehyde derivatives, -C(alkyl)=NH-NH-CONH2, an O-phenyl or -OCH2(phenyl) group, -C(cycloalkyl)=NOH, -C(cycloalkyl)=N-O(alkyl), an optionally substituted heterocycle; a heterocycle comprising a sulphur hetero atom; a cycloalkyl; a bicyclic group and preferably a norbomyl group; a phenyl ring fused to a heterocycle comprising a nitrogen hetero atom or to a carbocycle or a heterocycle bearing a keto function; an unbranched or branched lower alkyl comprising from 1 to 8 carbon atoms; an unbranched or branched alkyne comprising from 1 to 8 carbon atoms and preferably 1 to 5 carbon atoms; a linear or branched alkyl mono- or polysubstituted with phenyl groups which are either unsubstituted or monoor polysubstituted; a phenyl alkyl ketone in which the alkyl group is branched or unbranched or cyclic; a substituted or unsubstituted benzophenone; a substituted or unsubstituted, unbranched or branched or cyclic phenyl alcohol; an unbranched or branched alkene; a piperidyl group; a phenylcycloalkyl group; a polycyclic group, in particular a fluorenyl group, a naphthyl or polyhydronaphthyl group or an indanyl group; a phenol group; a ketone or keto derivative; a diphenyl group; a phenoxyphenyl group; a benzyloxyphenyl group.
- 31. Compound according to claim 29 or 30, characterized in that X^{II} is selected from -O-, -NH-, -CH₂-, -OCONH-, NHCO-, -NHCONH- and represents more preferably an oxygen atom.
- 32. Compound according to anyone of claims 29 to 31, characterized in that Y^{II} is selected from a linear or branched alkyl group; a cycloalkyl group, in particular cyclopentyl or cyclohexyl group; a phenyl group unsubstituted or monosubstituted, preferred substituent being halogen atom, in particular chorine; a heterocyclic radical, in particular pyridyl N-oxide or pyrazinyl radicals; a bicyclic radical such as a benzothiazolyl radical, Y^{II} being more preferably a phenyl group unsubstituted or mono-substituted as above-defined.
 - 33. Compound according to anyone of claims 29 to 31, characterized in that Y^{II} represents a phenyl group at least mono-substituted with a keto-substituent, in particular a linear or branched chain aliphatic ketone comprising from 1 to 8 carbon atoms and optionnally bearing a hydroxyl group, a cycloalkylketone, an aryl alkyl ketone or arylalkenylketone in which the aryl group is optionally substituted, or a heteroaryl ketone, preferably a cycloalkylketone; an oxime-substituent or an halogen atom.
- 34. Compound according to anyone of claims 29 to 31, characterized in that Y^{II} is a phenyl group at least mono-substituted with -CHO, a ketone, an aldehyde, -CH=CH-CHO, -C(alkyl)=N-OH, -C(alkyl)=N-O(alkyl) and other keto derivatives, -CH=N-OH, -CH=NO(alkyl) and other aldehyde derivatives, -C(cycloalkyl)=NOH, -C(cycloalkyl)=N-O(alkyl).
- 35. Compound according to anyone of claims 29 to 34, characterized in that chain A^{II} is a chain -(CH₂)_{nII}- with n varying from 1 to 6, preferably from 1 to 4, the chain A^{II} representing especially -(CH₂)₃-.
- 36. Compound according to anyone of claims 29 to 35, characterized in that the chain Bil is -(CH₂)₂- or -(CH₂)₃-.
- 37. Compounds according to anyone of claims 29 to 36, characterized in that X is an oxygen atom, the chain A represents -(CH₂)₃- and, for compounds of formula (IIa), the chain B represents -(CH₂)₃- also.
- 38. Compound according to anyone of claims 29 to 37, characterized in that it is one of the following compounds:
 - 3,3-Dimethylbutyl 3-piperidinopropyl ether
 - 3-Phenylpropyl 3-piperidinopropyl ether
 - 3-(4-Chlorophenyl)propyl 3-piperidinopropyl ether
 - 2-Benzothiazolyl 3-piperidinopropyl ether
 - N-Phenyl-3-piperidinopropyl carbamate
 - N-Pentyl-3-piperidinopropyl carbamate

- (S)-(+)-N-[2-(3,3-Dimethyl)butyl]-3-piperidinopropyl carbamate
- 3-Cyclopentyl-N-(3-(1-pyrrolidinyl)propyl)propanamide
- N-Cyclohexyl-N'-(1-pyrrolidinyl-3-propyl)urea
- 2-((2-Piperidinoethyl)amino)benzothiazole
- 5-Piperidinopentylamine

5

10

15

20

25

30

35

40

45

50

55

- 2-Nitro-5-(6-piperidinohexyl)pyridine
- 3-Nitro-2-(6-piperidinohexylamino)pyridine
- 2-(6-Piperidinohexylamino)pyrimidine
- N-(6-Phenylhexyl)piperidine
- N-phenyl-N'-(diethylamino-3-propyl)urea
- N-benzyl-N'-(3-piperidinopropyl)guanidine.
- 39. Compounds according to anyone of claims 1 to 15, having the following formula (III)

 $R^{\parallel 3} = \frac{1}{3} \times \mathbb{R}^{1}$ (III)

in which:

- NR¹R² is either in 3-position or in 4-position on the piperidyl moiety, R¹ and R² being as defined with reference to formula (A) in claim 1;
- R₂^{1II} denotes a linear or branched alkyl group having 1 to 6 carbon atoms; a piperonyl group, a 3-(1-benzimi-dazolonyl)propyl group; a group of formula

in which n_{III} is 0, 1, 2 or 3, X^{III} is a single bond or alternatively -O-, -S-, -NH-, -CO-, -CH=CH- or

R₃"

and R₃^{III} is H, CH₃, halogen, CN, CF₃ or an acyl group -COR₄^{III}, R₄^{III} being a linear or branched alkyl group having 1 to 6 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms or a phenyl group which can bear a CH₃ or F substituent; or alternatively a group of formula

` ---C--NH---R₅" Z"

in which Z^{III} denotes an O or S atom or a divalent group NH, N-CH₃ or N-CN and R₅^{III} denotes a linear or branched alkyl group having 1 to 8 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms which can

bear a phenyl substituent, a (C₃-C₆ cycloalkyl) (linear or branched, C₁-C₃ alkyl) group, a phenyl group which can bear a CH3, halogen or CF3 substituent, a phenyl(linear or branched, C1-C3 alkyl) group or a naphthyl, adamantyl or p-toluenesulphonyl group.

40. Compound according to claim 39, characterized in that R^{III} represents the group

Z^{III} and R^{III}₅ being as defined in claim 39, Z^{III} being especially O, S or NH.

- 41. Compound according to claim 40, characterized in that RIII₅ is a (C₃-C₆)cycloalkyl group.
- 42. Compound according to anyone of claims 39 to 41, which is N'-Cyclohexylthiocarbamoyl-N-1,4'-bipiperidine.
- 43. Compound according to anyone of claims 1 to 15, which have the following formula (IV):

$$\mathbb{R}^{\mathbb{N}}$$
 $\mathbb{R}^{\mathbb{N}}$ $\mathbb{R}^{\mathbb{N}}$ $\mathbb{R}^{\mathbb{N}}$

in which

10

15

20

25

30

35

40

45

50

- R¹ and R² are as defined with reference to general formula (A) in claim 1;
 R^{IV} represents a hydrogen atom or a group COR₃^{IV}, in which R₃^{IV} represents
- - (a) a linear or branched aliphatic group containing 1 to 11, and in particular 1 to 9, carbon atoms;
 - (b) a cyclane ring-system such as cyclopropane, phenylcyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, norbornane, adamantane, noradamantane, chlorooxonorbornane, chloroethylenedioxynorbornane, bromoethylenedioxynorbornane and the anhydride group of hydroxycarboxy-1,2,2trimethylcyclopentanecarboxylic acid;
 - (c) a benzene ring, unsubstituted or substituted at the para-position with a linear or branched aliphatic group containing 3 to 5 carbon atoms, as well as with a halogen;
 - (d) a group (CH₂)_{mIV}R₄IV in which m_{IV} is a number between 1 and 10, and R₄IV represents a cyclane ring system such as cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexane, norbornane, noradamantane, adamantane and 6,6-dimethylbicyclo[3.1.1] heptene; a benzene ring, unsubstituted or monosubstituted with a fluorine atom, a chlorine atom, a methyl group or a methoxy group; a thiophene ring grafted via its ring-position 2 or its ring-position 3; a carboxylic acid ester group COOR₅^{IV} in which R_5^{IV} is a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbornane; a carboxylic acid amide group of structure CONHR $_6^{IV}$, in which R_6^{IV} represents a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbornane; a carboxylic acid amide group of structure



in which the group

5

represents pyrrolidine, piperidine or 2,6-dimethylmorpholine; or an ether group - O-R₇^{IV}, it being possible for R₇^{IV} to be a benzene ring, unsubstituted or monosubstituted with a chlorine or fluorine atom or disubstituted with a chlorine atom and with a methyl group;

10

(e) a group -CH=CHR₈^{IV}, in which R₈^{IV} represents a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane, norbomane or norbornene; (f) a secondary amine group -NH(CH₂)_{nIV}R₉^{IV}, in which n_{IV} is a number between 1 and 5 and R₉^{IV} consti-

(f) a secondary amine group -NH(CH₂)_{nIV}R₉I^v, in which n_{IV} is a number between 1 and 5 and R₉I^v constitutes a cyclane ring-system such as cyclopropane, cyclobutane, cyclopentane, cyclohexane or norbornane, or a benzene ring, unsubstituted, mono-substituted with a fluorine or chlorine atom or with a methoxy group or trisubstituted with methoxy groups;

15

RIV also represents a hydroxyalkenyl group

. н

HO CH(CH₂)p_{IV}R₁₀IV

20

in which p_{IV} is a number between 2 and 9 and $R_{10}^{\ IV}$, represents a benzene ring or a phenoxy group; as well as a group

25

in which n_{IV} is a number between 1 and 5 and R_9^{IV} has the meaning stated above.

- 44. Compound according to claim 43, characterized in that R^{IV} represents the group COR₃^{IV}, R₃^{IV} representing especially an aliphatic group a).
 - 45. Compound according to anyone of claims 43 and 44, which is N-Heptanoyl-1,4'-bipiperidine.
- 35 46. Compound according to anyone of claims 1 to 15, having the following formula (VI):

40

45

50

55

$$R^{2} = R^{2} \times (CH_{2})_{mvi} \times (VI)$$

$$R^{2} \times (CH_{2})_{nvi} \times (VI)$$

wherein:

- A^{VI} is selected from -O-CO-NR $^1_{VI}$ -, -O-CO-, -NR $^1_{VI}$ -CO-NR $^1_{VI}$ -, -NR $^1_{VI}$ -CO-, -NR $^1_{VI}$ -, -O-, -CO-NR $^1_{VI}$ -, -CO-O-, and -C(=NR $^1_{VI}$)-NR $^1_{VI}$ -;
- the groups R¹_{VI}, which may be the same or different when there are two or three such groups in the molecule of formula VI, are selected from hydrogen, and lower alkyl, aryl, cycloalkyl, heterocyclic and heterocyclylalkyl groups, and groups of the formula -(CH₂)_{VVI}-G^{VI}, where G^{VI} is selected from CO₂R³_{VI}, COR³_{VI}, COR³_{VI}, COR³_{VI}, NR³_{VI}, NR³_{VI}, heteroaryl and phenyl, which phenyl is optionally substituted by halogen, lower alkoxy or polyhaloloweralkyl, and y_{VI} is an integer from 1 to 3;
- R²_{VI} is selected from hydrogen and halogen atoms, and alkyl, alkenyl, alkynyl and trifluoromethyl groups, and groups of the formula OR³_{VI}, SR³_{VI} and NR³_{VI};
- R³_{VI} and R⁴_{VI} are independently selected from hydrogen, and lower alkyl and cycloalkyl groups, or R³_{VI} and R⁴_{VI} together with the intervening nitrogen atom can form a saturated ring containing 4 to 6 carbon atoms that can be substituted with one or two lower alkyl groups;

- the group - $(CH_2)_{nVI}$ - A^{VI} - R^1_{VI} is at the 3- or 4-position, and the group R^2_{VI} is at any free position;
- m_{VI} is an integer from 1 to 3;
- and n_{VI} is 0 or an integer from 1 to 3.
- 5 47. Compound according to anyone of claims 1 to 15, having the following formula (VI):

$$R^{2} \qquad \qquad R^{2} \qquad \qquad R^{2} \qquad \qquad R^{2} \qquad \qquad (CH_{2})_{\Pi V I} \qquad A^{V I} \qquad (VI)$$

- wherein R¹_{VI} is an aryl group, preferably a phenyl group optionally substituted with a keto-substituent, in particular a linear or branched chain aliphatic ketone comprising from 1 to 8 carbon atoms and optionnally bearing a hydroxyl group, a cycloalkylketone, an aryl alkyl ketone or arylalkenylketone in which the aryl group is optionally substituted, or a heteroaryl ketone, preferably a cycloalkylketone, R²_{VI}, n_{VI}, m_{VI} and A^{VI} being as defined in claim 46.
- 48. Compound according to claim 46 or 47, characterized in that n_{VI} and m_{VI} are each 1, and A^{VI} represents an oxygen atom.
 - **49.** Compound according to claim 46 or 48, characterized in that R¹_{VI} is an aryl or -(CH₂)_{yVI}-G^{VI} with G^{VI} being a phenyl.
 - 50. Compound according to anyone of claims 46 to 49, which is one of the following compounds:
 - α-(4-Acetylphenoxy)-α'-piperidino p-xylol
 - α-(4-Acetylphenoxy)-α'-(1-pyrrolidinyl) p-xylol
 - α-(3-Phenylpropoxy)-α'-piperidino p-xylol.
 - 51. Compound according to anyone of claims 1 to 15, having the following formula (VII):

$$R^{1} \longrightarrow (CH_{2})n_{VII} \longrightarrow (CH_{2})_{MVII} \longrightarrow (VII)$$

in which

25

30

35

40

45

50

- R¹ and R² are as defined in reference to formula (A) in claim 1;
- X^{VII}, Y^{VII} and Z^{VII} are identical or different and represent O, N or S;
- n_{Vii} is varying from 1 to 3;
- m_{VII} is 1 or 2.
- **52.** Compound according to claim 51, characterized in that X^{VII} is 0 and Y^{VII} and Z^{VII} are each N to represent a 1, 2, 4-oxadiazolyl group.
- 53. Compound according to claims 51 or 52, which is one of the following compounds:
 - 3-(4-Chlorobenzyl)-5-(2-piperidinoethyl)-1,2,4-oxadiazole
 - compound 102.

54. Compound according to anyone of claims 1 to 15, having the following formula (VIII):

R¹ A^{VIII} B^{VIII} R²/VIII (VIII)

wherein ${\sf R}^1$ and ${\sf R}^2$ are as defined with reference to formula (A) in claim 1 and wherein ${\sf A}^{\sf VIII}$ is

1) a group of the formula $(CH_2)_{mVIII}$, wherein $m_{VIII} = 0-9$; or

2) a group of the formula:

5

10

15

20

25

30

35

40

45

50

55

wherein R^5_{VIII} represents hydrogen, (C_1-C_3) alkyl-, aryl (C_1-C_3) alkyl-, aryl-, wherein aryl may optionally be substituted, hydroxyl-, (C_1-C_3) alkoxy-, halogen, amino-, cyano- or nitro; and R^6_{VIII} represents hydrogen, (C_1-C_3) alkyl-, aryl (C_1-C_3) alkyl-, or aryl-, wherein aryl may optionally be substituted; or 3) a group of the formula:

wherein R^5_{VIII} and R^6_{VIII} are as defined above; or 4) a group of the formula:

if B^{VIII} is a group of the formula:

such that A^{VIII} and B^{VIII} together form a group of the formula:

wherein R^6_{VIII} is as defined above; or 5) a group of the formula:

wherein R^6_{VIII} is as defined above; or 6) a group of the formula:

if B^{VIII} is a group of the formula:

such that \mathbf{A}^{VIII} and \mathbf{B}^{VIII} together form a group of the formula:

wherein R^6_{VIII} is as defined above; or 7) a group of the formula:

wherein
$$x_{VIII} + y_{VIII} = m_{VIII}$$
-1; B^{VIII} is

1) a group of the formula:

5

10

wherein R⁵_{VIII} is as defined above; or 2) a group of the formula:

-c(

15

if A is a group of one of the formulas:

20

25

such that A and B together form a group of one of the formulas:

30

35

wherein R^6_{VIII} is as defined above; or 3) a group of the formula:

40

.

if X^{VIII} is a group of the formula:

45

50

such that \mathbf{B}^{VIII} and \mathbf{X}^{VIII} together form a group of the formula

55

wherein $p_{VIII} = 1-3$; or X^{VIII} is

- 1) a group of the formula $(CH_2)_{nVIII}$ wherein $n_{VIII} = 2-4$; or
- 2) a group of the formula:

if B^{VIII} is a group of the formula:

such that XVIII and BVIII together form a group of the formula:

wherein $p_{VIII} = 1-3$; or

3) two hydrogens (one on the carbon and one on the nitrogen); or

4) one hydrogen on the carbon atom and one R⁷_{VIII} group on the nitrogen atom,

wherein R⁷_{VIII} represents hydrogen, (C₁-C₁₀)alkyl-, aryl (C₁-C₁₀)alkyl-, or aryl, wherein aryl may optionally be substituted;

 Y^{VIII} is a group of the formula $(CH_2)_{kVIII}$, wherein $k_{VIII} = 0-2$;

 R^4_{VIII} represents hydrogen, (C_1 - C_{10})alkyl-, (C_1 - C_3)alkyl-sulfonamide-, aryl(C_1 - C_{10})alkyl-, aryl, wherein aryl may optionally be substituted;

or a group of the formula:

or a group of the formula:

wherein XVIII represents O, S, or NH,

R⁷VIII is as defined as above;

 $R_{V|II}^{8}$ represents (C₁-C₁₀)alkyl-, aryl(C₁-C₁₀)alkyl- or aryl, wherein aryl may optionally be substituted and wherein aryl is phenyl, substituted phenyl, naphtyl, substituted naphtyl, pyridyl;

55. Compound according to claim 54, having the formula

55

5

10

15

20

25

30

35

40

45

$$R^1$$
 $(CH_2)_{\Pi V | II}$ NH_2 R^2 $(CH_2)_{\Pi V | II}$ NH C NH $RV | R^2$ $(VIIIa)$ or R^2 $(VIIIb)$

R¹ and R² having the meaning given in claim 1 and n_{VIII} and R^{VIII} having the meaning given in claim 54.

- 56. Compound according to claim 54 or 55, which is 2-Nitro-5-(6-piperidinohexyl)pyridine.
- 57. Compound according to anyone of claims 1 to 15, having the following formula (IX):

wherein:

5

10

15

20

25

30

35

40

45

R¹ and R² are as defined with reference to formula (A) in claim 1.

 R_{IX}^1 is C_4 to C_{20} hydrocarbyl (in which one or more hydrogen atoms may be replaced by halogen, and up to four carbon atoms [and especially from 0 to 3 carbon atoms] may be replaced by oxygen, nitrogen or sulphur atoms, provided that R_{IX}^1 does not contain an -O-O-group),

 R^2_{IX} is H or C_1 to C_{15} hydrocarbyl (in which one or more hydrogen atoms may be replaced by halogen, and up to three carbon atoms may be replaced by oxygen, nitrogen or sulphur atoms, provided that R^2_{IX} does not contain an -O-O-group. m_{IX} is from 1 to 15 (preferably 1 to 10, more preferably 3 to 10, eg. 4 to 9) each X^{IX} group is independently

or one X^{IX} group is -N(R^4_{IX})-, -O- or -S- (provided that this X^{IX} group is not adjacent the -NR $^2_{IX}$ - group) and the remaining X^{IX} groups are independently

wherein $R^3_{|X}$ is H, C_1 to C_6 alkyl, C_2 to C_6 alkenyl, $-CO_2R^5_{|X}$, $-CON(R^5_{|X})_2$, $-CR^5_{|X2}OR^6_{|X}$ or $-OR^5_{|X}$ (in which $R^5_{|X}$ and $R^6_{|X}$ are H or C_1 to C_3 alkyl), and $R^4_{|X}$ is H or C_1 to C_6 alkyl.

- 50 58. Compound according to claim 57, which is compound 100.
 - 59. Compound according to anyone of claims 1 to 15, having the following formula (X):

wherein:

10

15

5


- R¹ and R² are as defined with reference to formula (A) in claim 1;
- R¹_x is H or CH₃;
- R²_x is selected from a phenyl optionally substituted with a halogen atom, preferably chlorine, a (C₁-C₄)alkyl, a (C₁-C₄)alkoxy, CF₃, OCF₃, NO₂, NH₂; or a CH₂-phenyl optionally substituted as above-specified;
- 60. Compound according to claim 59, which is compound 101.
- 61. Compound according to claims 1 to 15, having the following formula (XI):

25

35

40

20



where R1 and R2 are as defined with reference to formula (A) in claim 1; 30

where AXI is -NHCO-, -N(CH3)-CO-, -NHCH2-, -N(CH3)-CH2-, -CH=CH-, -COCH2-, CH2CH2-, -CH(OH)CH2-, or -C=C-:


XXI is H, CH₃, NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, or SH; R₂XI is hydrogen or a methyl or ethyl group; R₃XI is hydrogen or a methyl or ethyl group; nXI is 0, 1, 2, 3, 4, 5 or 6; and

R₁XI is selected from the group consisting of C₃ to C₈ cycloalkyl; phenyl or substituted phenyl; decahydronaphthalene and octahydroindene; or

 R_1^{XI} and X^{XI} may be taken together to denote a 5,6- or 6,6-saturated bicyclic ring structure when X^{XI} is NH, O, S, or SO₂.

- 62. Compound according to claim 61, characterized in that it is one of the following compounds 104, 105 or 106.
- 63. Compound according to claim 1 to 15, having the following formula (XII):

45



50

55

where R1 and R2 are as defined in reference to formula (A) in claim 1;

where R₂^{XII} is a hydrogen or a methyl or ethyl group;

R₃^{XII} is a hydrogen or a methyl or ethyl group; n^{XII} is 0, 1, 2, 3, 4, 5, or 6; and

R₁XII is selected from the group consisting of C₃ to C₈ cycloalkyl; phenyl or substituted phenyl; alkyl; heterocyclic;

decahydronaphthalene; and octahydroindene;

with the provisos that

5

10

30

35

40

45

when XXII is H, AXII can be -CH2CH2-, -COCH2-, -CONH-, -CON(CH3)-, -CH=CH-, -C=C-, -CH2-NH-, -CH2-N(CH₃)-, -CH(OH)CH₂-, -NH-CH₂-, -N(CH₃)-CH₂-, -CH₂O-, -CH₂S-, or -NHCOO-;

when X^{XII} is NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, CH₃, SH or SCH₃; A^{XII} can be -NHCO-, -N(CH₃)-CO-, -NHCH₂-, -N(CH₃)-CH₂-, -CH=CH-, -COCH₂-, -CH₂CH₂-, -CH(OH)CH₂-, or —C=C-; and when R₁^{XII} and X^{XII} taken together denote a 5,6 or 6,6 saturated bicyclic ring structure X^{XII} can be NH, O, or S.

- 64. Compound according to claim 63, characterized in that, AXII is —CH=CH- or —C=C-.
- 65. Compound according to claims 63 to 64, characterized in that R₂^{XII}, R₃^{XII} are each hydrogen atom.
- 66. Compound according to anyone of claims 63 to 65, characterized in that n_{XII} is an alkyl group.
- 67. Compound according anyone of claims 63 to 66, which is compound 107.
 - 68. Compound according to anyone of claims 1 to 15 having the following formula (XIII):

wherein R^1 and R^2 are as defined with reference to formula (A) in claim 1. wherein D^{XIII} is CH_2 or CH_2 - CH_2 , Z^{XIII} represents sulfur (S) or oxygen (O), preferably O, X_{XIII} is 0 or 1, n_{XIII} is an integer from 0 to 6,

and R₂XIII represents a substituted or unsubstituted linear chain or branched chain alkyl group of up to about 20 carbon atoms, a substituted or unsubstituted carbocyclic group of up to about 20 carbon atoms including mono and bicyclic moieties, and a substituted or an unsubstituted aryl group of up to about 20 carbon atoms, or any combination of above-mentioned groups, or salts thereof.

- 69. Compound according to claim 68, which is compound 103.
- 70. Compound according to anyone of claims 1 to 15, having the following formula (XIV)

$$\begin{array}{c|c}
R^{1}_{XIV} & R^{2}_{XIV} \\
R^{1} & & & \\
R^{2} & & & \\
R^{2} & & & \\
\end{array}$$

$$\begin{array}{c|c}
R^{2}_{XIV} & & & \\
\hline
TXIV & N - R^{5}_{XIV} \\
\hline
(C)_{D_{YN}} & R^{4}_{XIV}
\end{array}$$
(XIV)

wherein R1 and R2 are as defined in reference of formula (A) in claim 1;

- (4) -C(O)OR^{7′}_{XIV}; wherein R^{7′}_{XIV} is the same as R⁷_{XIV} defined below except that R^{7′}_{XIV} is not H;
- (5) -C(O)R⁷_{XIV};
 - (6) -C(O)NR⁷XIVR⁸XIV;
 - (7) allyl;
 - (8) propargyl; and

50

- (9) -(CH₂)_q-R⁶_{XIV} wherein q_{XIV} and R⁶_{XIV} are as defined above, and when q_{XIV} is equal to 1, then R⁶_{XIV} is not OH or SH:
 - (E) $R_{X|V}^7$ and $R_{X|V}^8$ are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and C₃ to C₆ cycloalkyl;
 - (F) the dotted line (-----) represents a double bond that is optionally present when m_{XIV} is 1, and n_{XIV} is not 0, and p is not 0 (i.e., the nitrogen in the ring is not bound directly to the carbon atom bearing the double bond), and when said double bond is present then R^2_{XIV} is absent; and
 - (G) when m_{XIV} is 2, each R^1_{XIV} is the same or different substituent for each m_{XIV} , and each R^2_{XIV} is the same or different substituent for each m_{XIV} and at least two of the substituents R^1_{XIV} and/or R^2_{XIV} are H. (A) m_{XIV} is an integer selected from the group consisting of: 1 and 2;
 - (B) n_{XIV} and p_{XIV} are intergers and are each independently selected from the group consisting of: 0, 1, 2, 3, and 4 such that the sum of n_{XIV} and p_{XIV} is 4 and p_{XIV} is a 6-membered ring;
 - (C) R³_{XIV} and R⁴_{XIV} are each independently bound to the same or different carbon atom of ring T^{XIV}, such that there is only one R³_{XIV} group and one R⁴_{XIV} group in ring T^{XIV}, and each R¹_{XIV}, R²_{XIV}, R³_{XIV} and R⁴ is independently selected from the group consisting of:
 - (1) H;

5

10

15

20

25

30

35

40

45

50

- (2) C₁ to C₆ alkyl; and
- (3) -(CH₂)_{qXIV}-R⁶_{XIV} wherein q_{XIV} is an integer of: 1 to 7, and R⁶_{XIV} is selected from the group consisting of: phenyl, substituted phenyl, -OR⁷_{XIV}, -C(O)OR⁷_{XIV}, -C(O)R⁷_{XIV}, -OC(O)R⁷_{XIV}, -C(O)NR⁷_{XIV}, CN and -SR⁷_{XIV} wherein R⁷_{XIV} and R⁸_{XIV} are as defined below, and wherein the substituents on said substituted phenyl are each independently selected from the group consisting of: -OH, -O-(C₁ to C₆)alkyl, halogen, C₁ to C₆ alkyl, -CF₃, -CN, and -NO₂, and wherein said substituted phenyl contains from 1 to 3 substituents:
- (D) R⁵_{XIV} is selected from the group consisting of:
 - (1) H:
 - (2) C₁ to C₂₀ alkyl;
 - (3) C₃ to C₆ cycloalkyl;
- 71. Compound according to claim 70, which is selected from compounds having the following formula (XIVa), (XIVb) or (XIVc)

in which R⁵_{XIV} is preferably H or CH₃ and R³_{XIV} and R⁴_{XIV} are preferably each H.

72. Compounds according to anyone of claims 1 to 15, having the following formula (XV):

$$\begin{array}{c|c}
R^{1}_{XV} & R^{2}_{XV} & R^{3}_{XV} & R^{4}_{XV} \\
R^{1}_{XV} & & & & & & & & \\
R^{1}_{XV} & & & & & & & \\
R^{2}_{XV} & & & & & & & \\
R^{2}_{XV} & & & & & & \\
R^{3}_{XV} & & & & & & \\
R^{6}_{XV} & & & & & \\
\end{array} (XV)$$

where R1 and R2 are as defined in reference to formula (A) in claim 1;

- (A) m_{XV} is an integer selected from the group consisting of: 0,1, and 2;
- (B) n_{XV} and p_{XV} are intergers and are each independently selected from the group consisting of: 0, 1, 2, and 3 such that the sum of n_{XV} and p_{XV} is 2 or 3 such that when the sum of n_{XV} and p_{XV} is a 4-membered ring and when the sum of n and p_{XV} is 3, T^{XV} is a 5-membered ring; (C) each R^1_{XV} R^2_{XV} R^3_{XV} R^6_{XV} R^6_{XV} R^7_{XV} and R^8_{XV} is independently selected from the group consisting of:
- (1) H;

25

35

40

45

50

- (2) C₁ to C₆ alkyl;
- (3) C₃ to C₆ cycloalkyl; and
- (4) -(CH₂)_{qXV}-R⁹_{XV} wherein q_{XV} is an integer of: 1 to 7, and R⁹_{XV} is selected from the group consisting of: phenyl, substituted phenyl, -OR¹⁰_{XV}, -C(O)OR¹⁰_{XV}, -C(O)R¹⁰_{XV}, -OC(O)R¹⁰_{XV}, -C(O)NR¹⁰_{XV}, CN and -SR¹⁰_{XV} wherein R¹⁰_{XV} and R¹¹_{XV} are as defined below, and wherein the substituents on said substituted phenyl are each independently selected from the group consisting of: -OH, -O-(C1 to C6) alkyl, halogen, C1 to C6 alkyl, -CF3, -CN, and -NO2, and wherein said substituted phenyl contains from 1 to 3 substituents; examples of -(CH₂)_{qXV}R⁹_{XV} include benzyl, substituted benzyl and the like, wherein the sub-

stitutents on the substituted benzyl are as defined above for said substituted phenyl;

- (D) R⁵_{XV} is selected from the group consisting of:
 - (1) H;

5

10

15

20

25

30

35

40

45

- (2) C₁ to C₂₀ alkyl;
- (3) C₃ to C₆ cycloalkyl;
- (4) -C(O)OR^{10′}_{XV}; wherein R^{10′}_{XV} is the same as R¹⁰_{XV} defined below except that R^{10′}_{XV} is not H; (5) -C(O)R¹⁰_{XV}; (6) -C(O)NR¹⁰_{XV}R¹¹_{XV};

- (7) allyl;
- (9) -(CH₂) $_{qXV}$ -R⁹ $_{XV}$, wherein q_{XV} and R⁹ $_{XV}$ are as defined above with the proviso that when q_{XV} is 1 then R9XV is not -OH or -SH;
- (E) $R^{10}{}_{XV}$ and $R^{11}{}_{XV}$ are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and C₃ to C₆ cycloalkyl; and, for the substituent -C(O)NR¹⁰ ${}_{XV}R^{11}$, $R^{10}{}_{XV}$ and $R^{11}{}_{XV}$ together with the nitrogen to which they are bound, can form a ring having 5, 6, or 7 atoms;
- (F) the dotted line (----) represents a double bond that is optionally present when m_{XV} is 1, and T^{XV} is a 5membered ring, and nXV is not 0, and pXV is not 0 (i.e., the nitrogen in the ring is not bound directly to the carbon atom bearing the double bond), and when said double bond is present then R^2_{XV} and R^8_{XV} are absent;
- (G) when m_{XV} is 2, each R^1_{XV} is the same or different substituent for each m_{XV} and each R^2_{XV} is the same or different substituent for each mxv;
- (H) when n_{XV} is 2 or 3, each R^3_{XV} is the same or different substituent for each n_{XV} and each R^4_{XV} is the same or different substituent for each nxv; and
- (I) when p_{XV} is 2 or 3, each R^6_{XV} is the same or different substituent for each p, and each R^7_{XV} is the same or different substituent for each pxv
- 73. Compound according to anyone of claims 1 to 15, having the following formula (XVI)

$$\begin{array}{c|c}
R^1 & NR_2^{XVI} \\
R^2 & NR_5^{XVI} & NR_5^{XVI} \\
\end{array}$$
(XVI)

where R1 and R2 are as defined in reference to formula (A) in claim 1; Z^{XVI} is a group of the formula $(CH_2)_{mXVI}$ wherein $m_{XVI} = 1-5$ or a group of the formula:

wherein

 $R_{XVI}^6 = (C_1 - C_3)alkyl$

 $R_{XVI}^{7} = (C_1 - C_3)$ alkyl; wherein Z_{XVI}^{7} may optionally comprise other substituents selected such that the activity of the derivative is not neg-50 atively affected,

XXVI represents S, NH or CH2

R¹_{XVI} represents hydrogen, (C₁-C₃)alkyl-, aryl(C₁-C₁₀)alkyl, wherein aryl may optionally be substituted, aryl, (C₅-C₇)cycloalkyl(C₁-C₁₀)alkyl-, or a group of the formula:

wherein $n_{XVI} = 1-4$, R^8_{XVI} is aryl, aryl(C_1-C_{10})alkyl-, (C_5-C_7)cycloalkyl- or (C_5-C_7) cycloalkyl(C_1-C_{10})alkyl-, and R^9_{XVI} is hydrogen, (C_1-C_{10})alkyl-or aryl; R_2^{XVI} and R_5^{XVI} represent hydrogen, (C_1-C_3)alkyl-, aryl or arylalkyl-, wherein aryl may optionally be substituted; wherein aryl is phenyl, substituted phenyl, naphthyl, substituted napththyl, pyridyl or substituted pyridyl.

74. Compound according to anyone of claims 1 to 15, having the following formula (XVII):

wherein m_{XVII} represents an integer of from 4 to 6.

R⁴_{XVII} represents a hydrogen atom, a linear or branched alkyl group, a cycloalkyl group, a cycloalkylalkyl group, a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group; and Z^{XVII} represents R⁵_{XVII} or A^{XVII}-R⁶_{XVII}, wherein A^{XVII} represents S or O, R₅^{XVII} represents a hydrogen atom, a lower alkyl group, a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group, and R₆^{XVII} represents a lower alkyl group, a lower alkenyl group, a lower alkynyl group or a substituted or unsubstituted aralkyl group.

75. Compound according to anyone of claims 1 to 15, having the following formula (Va) or (Vb):

or $R^{1} N - (CH_{2})_{q\overline{V}} Z^{V}$ (Vb)

50 in which

- R¹ and R² are as defined with reference to formula (A) in claim 1;
- Z^V represents NH, O or S;
- XV represents a hydrogen atom or a lower alkyl
- 55 q_V is 2 to 5.

5

10

15

20

40

76. Pharmaceutical composition characterized in that it comprises as active ingredient, a therapeutically effective amount of a compound according to anyone of claim 1 to 75 in combination with a pharmaceutically acceptable

vehicle or excipient.

5

15

20

25

30

35

40

45

50

55

- 77. Medicament acting as an antagonist and/or agonist of the histamine H₃-receptors, characterized in that it comprises as active ingredient, an effective amount of a compound according to anyone of claims 1 to 74.
- 78. Use of a compound having the following general formula (A): in which:

$$[W]-N \stackrel{R^1}{\underset{R^2}{\longleftarrow}}$$
 (A)


- W is a residue which imparts antagonistic and/or agonistic activity at histamine H₃-receptors when attached to an imidazole ring in 4(5)-position;
- R1 and R2 may be identical or different and represent each independently
 - · a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
 - · a saturated nitrogen-containing ring

with m ranging from 2 to 8, or

a non-aromatic unsaturated nitrogen-containing ring

with p and q being from 0 to 3 independently and r being from 0 to 4, provided that p and q are not simulteously 0 and $2 \le p + q + r \le 8$, R^{a-d} being independently a hydrogen atom or a lower alkyl, cycloalkyl, or carboalkoxy group, or

- · a morpholino group, or
- a N-substituted piperazino group:



with R being a lower alkyl, cycloalkyl, carboalkoxy, aryl, arylalkyl, an alkanoyl or aroyl group, as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, for the preparation of a medicament acting as an antagonist and/or agonist of the histamine H₃-receptors.

79. Use according to claim 78, characterized in that compound (A) is a compound of general formula (I)

$$(R^3)_{n_3}$$
 $X-C_nH_{2n}-N$ R^1 (I)

in which:

10

15

20

25

35

40

45

50

- C_nH_{2n} is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- R¹ and R² are as defined in claim 78;
- n₃ is an integer from 0 to 5;
- R³ represents each independently
 - · a halogen atom,
 - a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino, α-hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
 - or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring.
- 80. Use according to claim 78, characterized in that compound (A) is as defined in any one of claims 2 to 69.
- 81. Use according to claim 78 characterized in that compound (A) is one of the following compounds:
 - 1-(5-phenoxypentyl)-piperidine
 - 1-(5-phenoxypentyl)-pyrrolidine
- 30 N-methyl-N-(5-phenoxypentyl)-ethylamine
 - 1-(5-phenoxypentyl)-morpholine
 - N-(5-phenoxypentyl)-hexamethyleneimine
 - N-ethyl-N-(5-phenoxypentyl)-propylamine
 - 1-(5-phenoxypentyl)-2-methyl-piperidine
 - 1-(5-phenoxypentyl)-4-propyl-piperidine
 - 1-(5-phenoxypentyl)-4-methyl-piperidine
 - 1-(5-phenoxypentyl)-3-methyl-piperidine
 - 1-(3-prierioxyperityr)-3-metryr-piperidine
 - 1-acetyl-4-(5-phenoxypentyl)-piperazine
 - 1-(5-phenoxypentyl)-3,5-trans-dimethyl-piperidine
 - 1-(5-phenoxypentyl)-3,5-cis-dimethyl-piperidine
 - 1-(5-phenoxypentyl)-2,6-cis-dimethyl-piperidine
 - 4-carboethoxy-1-(5-phenoxypentyl)-piperidine
 - 3-carboethoxy-1-(5-phenoxypentyl)-piperidine
 - 1-(5-phenoxypentyl)-1,2,3,6-tetrahydropyridine
 - 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 - 1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
 - 1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
 - 1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
 - 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
 - 1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine
 - 1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine
 - 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
 - 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
 - 1-{5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
- 55 1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
 - 1-(5-phenoxypentyl)-2,5-dihydropyrrole
 - 1-{5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl}-pyrrolidine
 - 1-(4-phenoxybutyl)-pyrrolidine

	1-(6-phenoxyhexyl)-pyrrolidine
	1-(5-phenylthiopentyl)-pyrrolidine
	1-(4-phenylthiobutyl)-pyrrolidine
	1-(3-phenoxypropyl)-pyrrolidine
5	1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
J	1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
	1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
40	1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
10	
	N-[3-(4-nitrophenoxy)-propyl]-diethylamine
	N-[3-(4-cyanophenoxy)-propyl]-diethylamine
	1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
	1-{5-[4-(phenylacetyl)-phenoxy]-pentyl}-pyrrolidine
15	N-[3-(4-acetylphenoxy)-propyl]-diethylamine
	1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
	1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
20	1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
	1-[5-(4-cyanophenoxy)-pentyl]-piperidine
	N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
	N-[2-(4-cyanophenoxy)-ethyl]-diethylamine
	N-[3-(4-cyanophenoxy)-propyl]-dimethylamine
25	N-[4-(4-cyanophenoxy)-butyl]-diethylamine
	N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
	1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
	1-[3-(4-cyanophenoxy)-propyl]-piperidine
	N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
30	N-[6-(4-cyanophenoxy)-hexyl]-diethylamine
	N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
	N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
	4-(3-diethylaminopropoxy)-acetophenone-oxime
	1-[3-(4-acetylphenoxy)-propyl]-piperidine
35	1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
55	1-[3-(4-acetylphenoxy)-propyi]-3,5-trans-dimethyl-piperidine
	1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
	1-[3-(4-propionylphenoxy)-propyl]-piperidine
	1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
40	1-[3-(4-formylphenoxy)-propyl]-piperidine
	1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
	N-[3-(4-propionylphenoxy)-propyl]-diethylamine
	1-[3-(4-butyrylphenoxy)-propyl]-piperidine
	1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine
45	

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, for the preparation of a medicament acting as an antagonist of the histamine H₃-receptors.

82. Use according to claim 78, characterized in that compound (A) is one of the following compounds:

1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
N-[3-(4-cyanophenoxy)-propyl]-diethylamine
N-[3-(4-acetylphenoxy)-propyl]-diethylamine
55
1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine
N-[4-(4-cyanophenoxy)-butyl]-diethylamine
1-[3-(4-cyanophenoxy)-propyl]-piperidine
N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine

N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine

- 4-(3-diethylaminopropoxy)-acetophenone-oxime
- 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
- 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
- 1-[3-(4-propionylphenoxy)-propyl]-piperidine

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, for the preparation of a medicament acting as an antagonist of the histamine H₃-receptors.

- 83. Use according to claim 78, characterized in that compound (A) is one of the following compounds:
 - 3,3-Dimethylbutyl 3-piperidinopropyl ether,
 - 3-Phenylpropyl 3-piperidinopropyl ether,
- 15 3-(4-Chlorophenyl)propyl 3-piperidinopropyl ether,
 - 2-Benzothiazolyl 3-piperidinopropyl ether,
 - N-Phenyl-3-piperidinopropyl carbamate,
 - N-Pentyl-3-piperidinopropyl carbamate,
 - (S)-(+)-N-[2-(3,3-Dimethyl)butyl]-3-piperidinopropyl carbamate,
- 20 N-(4-Chlorobenzyl)-S-(3-piperidinopropyl) isothiourea,
 - N'-Cyclohexylthiocarbamoyl-N-1,4'-bipiperidine,
 - N-Heptanovl-1,4'-bipiperidine,
 - 3-Cyclopentyl-N-(3-(1-pyrrolidinyl)propyl)propanamide,
 - N-Cyclohexyl-N'-(1-pyrrolidinyl-3-propyl)urea,
 - α-(4-Acetylphenoxy)-α'-piperidino p-xylol,
 - α -(4-Acetylphenoxy)- α '-(1-pyrrolidinyl) p-xylol,
 - α -(3-Phenylpropoxy)- α '-piperidino p-xylol,
 - 3-(4-Chlorobenzyl)-5-(2-piperidinoethyl)-1,2,4-oxadiazole,
 - 2-((2-Piperidinoethyl)amino)benzothiazole,
- 30 5-Piperidinopentylamine,

5

10

25

40

- 2-Nitro-5-(6-piperidinohexyl)pyridine,
- 3-Nitro-2-(6-piperidinohexylamino)pyridine,
- 2-(6-Piperidinohexylamino)pyrimidine,
- N-(6-Phenylhexyl)piperidine,
- 35 compounds 98 to 107.

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, for the preparation of a medicament acting as an antagonist of the histamine H₃-receptors.

- **84.** Medicament according to anyone of claims 77 to 83, for the treatment of central nervous system disorders, in particular Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.
- 85. Medicament according to anyone of claims 77 to 83, having psychotropic effects, promoting wakefulness, attention, memory and improving mood, intended to be used in particular in the treatment of Alzheimer disease and other cognitive disorders in aged persons, depressive or asthenic states.
- **86.** Medicament according to anyone of claims 77 to 82, having nootropic effects, intended to be used in particular in treatment to stimulate attention and memorization capacity.
 - 87. Medicament according to anyone of claims 77 to 83, for the treatment of obesity, vertigo and motion sickness.
- **88.** Medicament according to anyone of claims 77 to 83, for the treatment of CNS disorders, in particular of aged persons.

Europäisches Patentamt

European Patent Office

Office européen des brevets



EP 0 982 300 A3 (11)

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3: 08.03.2000 Bulletin 2000/10

(43) Date of publication A2: 01.03.2000 Bulletin 2000/09

(21) Application number: 98403351.4

(22) Date of filing: 31.12.1998

(51) Int. Cl.⁷: **C07D 295/088**, C07C 211/08, C07D 211/04, C07D 295/185, C07D 211/62, C07D 211/70, C07D 207/20, A61P 25/28

(84) Designated Contracting States: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

Designated Extension States: AL LT LV MK RO SI

(30) Priority: 29.07.1998 EP 98401944

(71) Applicant: SOCIETE CIVILE BIOPROJET F-75003 Paris (FR)

(72) Inventors:

- Schwartz, Jean-Charles 75014 Paris (FR)
- · Arrang, Jean-Michel 91410 Dourdan (FR)
- · Garbarg, Monique 75017 Paris (FR)

- Lecomte, Jeanne-Marie 75003 Paris (FR)
- · Ligneau, Xavier 75013 Paris (FR)
- · Schunack, Walter G. 14129 Berlin (DE)
- · Stark, Holger 14199 Berlin (DE)
- · Ganellin, Charon Robin Welwyn, Hert AL6 0TD (GB)
- · Leurquin, Fabien London E2 6DZ (GB)
- · Sigurd, Elz 12107 Berlin (DE)

(74) Representative:

Le Guen, Gérard et al **CABINET LAVOIX** 2, place d'Estienne d'Orves 75441 Paris Cédex 09 (FR)

Non-imidazole alkylamines as histamine H3 - receptor ligands and their therapeutic (54)applications

(57)Compounds of formula (A):

$$[W]-N <_{R^2}^{R^1}$$
 (A)

wherein:

- W is a residue which imparts antagonistic and/or agonistic activity at histamine H3-receptors when attached to an imidazole ring in 4(5) position.
- R1 and R2 may be identical or different and represent each independently
 - a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached.
 - a saturated nitrogen-containing ring (i) as defined,

- a non-aromatic unsaturated nitrogen-containing ring (ii) as defined,
- a morpholino group, or
- a N-substituted piperazino group as defined.

Compounds (A) are useful for preparing medicaments acting as antagonists and/or agonists at the H₃receptors of histamine.



EUROPEAN SEARCH REPORT

Application Number EP 98 40 3351

Category	Citation of document with in- of relevant passa		Relev to clai		CLASSIFICATION OF THE APPLICATION (Int.CL7)
X	Quaternary Ammonium Aliphatic Tertiary J.AMER.CHEM.SOC., vol. 52, 1930, page	MARVEL C. S.: "Cyclic Salts from Halogenated Amines" s 287-294, XP002084866 90; claim 1; examples	16-28	3	·
X	J.MED.CHEM., vol. 24, no. 2, 198 XP000565653	vity of Substituted benzene Derivatives"	16-28	3,79	
X	and Related Compoun J.MED.CHEM., vol. 14, no. 9, 197 XP002084867	N-Arylethylenediamines ds as Anorectic Agents"	16-28	8	TECHNICAL FIELDS SEARCHED (Im.Cl.7)
X	yl]piperidines and Compounds" J.MED.CHEM., vol. 32, no. 1, 198 XP002084868	y of hyl)-1-[3-(aryloxy)prop Structurally Related 9, pages 105-118, 90; claim 1; examples	16-2	8	
	The present search report has	been drawn up for all claims			
	Place of search	Date of completion of the search	' 		Examiner
	MUNICH	14 October 1999		Scr	uton-Evans, I
X:pa Y:pa do A:te	CATEGORY OF CITED DOCUMENTS rticularly relevant if taken alone rticularly relevant if combined with anot current of the same category chnological background on-written disclosure termediate document	T : theory or principl E : earlier patent do after the filing dat	curnent, but te in the appli or other re	ng the it ut publis lication easons	nvention shed on, or

3

מונד בי הייור בי בי הייור אים



EUROPEAN SEARCH REPORT

Application Number EP 98 40 3351

Category	Citation of document with ind		Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.7)
X	of relevant passage KIKUMOTO R ET AL: AGGREGATION INHIBITO PROPERTIES OF 2- (OMEGA-AMINOALKOXY) F JOURNAL OF MEDICINAL vol. 33, no. 6, June 1818-1823, XP0006734 * page 334; claim 1; tables I-III *	SYNTHESES AND PLATELET RY AND ANTITHROMBOTIC PHENYLETHYLBENZENES" CHEMISTRY, 1990, pages		APPECATION (Inco.)
X	CHENEY L.C. ET AL.: Ethers of the Benzy J.AMER.CHEM.SOC., vol: 71, 1949, page: * page 60; claim 1; I *	"Alkylaminoalkyl Iphenols" s 60-64, XP002086293 examples 1-5,7; table	16-28,79	
A STARK H. ET AL.: Histamine H3-rece DRUGS OF THE FUTU vol. 21, no. 5, 1 XP002084872		or Antagonists"	16-28,79	TECHNICAL FIELDS SEARCHED (Int.Ct.7)
	The present search report has			
	Place of search	Date of completion of the search	·	Examiner
Y:p da A:te	MUNICH CATEGORY OF CITED DOCUMENTS articularly relevant if taken alone articularly relevant if combined with anot sourment of the same category schnological background on-written disclosure termediate document	T: theory or principle: E: earlier patent of after the filing of the cournent clade L: document clade A: member of the document	ote underlying the ocument, but published in the application for other reasons	ished on, or



LACK OF UNITY OF INVENTION SHEET B

Application Number

EP 98 40 3351

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

8. Claims: 61-67

A compound of the formula XI or XII wherein the only comon feature with subjects 1-7 is NR1R2, and which has a cyclic moiety Rlxi or Rlxii

9. Claims: 70-72

Compounds with 4,5,or 6-membered N-containing ring, the only comon feature with subjects 1-8 being the NR1R2 group.

10. Claim: 74

Compounds of formula XVII, characterised in the presence of an N-containing heterocycle, which is N-substituted by C=N-R4, and directly substituted by the NR1R2 group, the only common group with subjects 1-9

11. Claim: 75

Compound swith a thiazole or benzthiazole ring, 2-substituted by the NR1R2-linker, with only the NR1R2 in common with the subjects 1-10

12. Claims: 81-83

Use of specifically named compounds, whose only common feature is a NR1R2 group, and which form no unifying common concept, as they have all such diverse structures

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 40 3351

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-10-1999

cited in search	ment report	Publication date	Patent family member(s)	Publication date
US 394743	4 A		NL 7404135 A SE 391925 B SE 7405652 A ZA 7400683 A	10-12-197 07-03-197 09-12-197 24-09-197
US 475130	2 A	14-06-1988	JP 58159471 A JP 58159472 A JP 58159473 A JP 58159474 A JP 58159475 A JP 58159476 A JP 58159477 A EP 0090972 A US 4533731 A	21-09-198 21-09-198 21-09-198 21-09-198 21-09-198 21-09-198 21-09-198 06-08-198
GB 924961	А		BE 588558 A CH 361005 A CH 395126 A CH 436259 A DE 1238485 B FR 558 M FR 84256 E FR 1421206 A GB 824853 A GB 921978 A LU 38374 A NL 129619 C NL 249341 A	05-05-190 09-03-190
US 331269	6 A	04-04-1967	NONE	
DE 262426	1 A	16-12-1976	FR 2313042 A FR 2349332 A BE 842453 A CH 597192 A GB 1513092 A	31-12-19 25-11-19 01-10-19 31-03-19 07-06-19
US 287015	1 A		NONE	
	С		NONE	

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

9

BNSDOCID: <EP_ ____0982300A3_I_>

£.

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
Потиев.

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.